


# Researchers propose social network modeling to fight hospital infections

22 October 2013



This is an illustration of dense (left) and sparse (right) patient ICU social networks. Patients in the circle that share a nurse are connected by lines (links), while patients that share a physician have the same shade/color (gray or black). Credit: University of Maryland

Two researchers at the University of Maryland's Robert H. Smith School of Business have teamed up with a researcher at American University to develop a framework to help prevent costly and deadly infections acquired by hospitalized patients. According to the Department of Health and Human Services (HHS), these transmissions strike one out of every 20 inpatients, drain billions of dollars from the national health care system and cause tens of thousands of deaths annually.

The research of Sean Barnes, Smith School assistant professor of operations management; Bruce Golden, the Smith School's France-Merrick Chair in Management Science; and Edward Wasil of American's Kogod School of Business, utilized computer models that simulate the interactions between patients and [health care workers](#) to determine if these interactions are a source for

spreading multi-drug resistant organisms (MDROs). Their study shows a correlation of a "sparse, social [network structure](#)" with low infection transmission rates.

This study comes in advance of HHS' 2015 launch and enforcement of a new initiative that penalizes hospitals at an estimated average rate of \$208,642 for violating specific requirements for infection control. In response, the study's authors have introduced a conceptual framework for hospitals to model their social networks to predict and minimize the spread of bacterial infections that often are resistant to antibiotic treatments.

The authors manipulated and tracked the dynamics of the social network in a mid-Atlantic hospital's intensive care unit. They focused on interactions between patients and health care workers – primarily nurses – and the multiple competing factors that can affect transmission.

"The basic reality is that healthcare workers frequently cover for one another due to meetings, breaks and sick leave," said Barnes. "These factors, along with the operating health care-worker-to-patient ratios and patient lengths of stay, can significantly affect transmission in an ICU... But they also can be better controlled."

The next step is to enable hospitals to adapt this framework, which is based on maximizing staff-to-patient ratio to ensure fewer nurses and physicians come in contact with each patient, especially high-risk patients.

"The [health care](#) industry's electronic records movement could soon generate data that captures the structure of patient-healthcare worker interaction in addition to multiple competing, related factors that can affect MDRO [transmission](#)," said Barnes.

The study, "Exploring the Effects of Network

Structure and Healthcare Worker Behavior on the Transmission of Hospital-Acquired Infections," appears in a recent issue of the peer-reviewed *IIE Transactions on Healthcare Systems Engineering*. The study was partially funded by the Robert H. Smith School of Business Center for Health Information and Decision Systems.

**More information:** A full copy of the study is available at: [www.tandfonline.com/doi/abs/10 ... =uhse20#.UmV9WPmsjIN](http://www.tandfonline.com/doi/abs/10.1080/17513758.2013.828282)

Provided by University of Maryland

APA citation: Researchers propose social network modeling to fight hospital infections (2013, October 22) retrieved 20 January 2022 from <https://medicalxpress.com/news/2013-10-social-network-hospital-infections.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*