

Less 'brown fat' could help explain why a fifth of the world's population is highly susceptible to type 2 diabetes

11 November 2013

Lower amounts of brown adipose tissue (BAT, or 'brown fat') could help explain why south Asians—who make up a fifth of the world's population—have an exceptionally high susceptibility to developing metabolic problems such as obesity and type 2 diabetes. The findings, published in *The Lancet Diabetes & Endocrinology*, could lead to new ways of preventing diabetes.

BAT has shown beneficial effects on glucose tolerance, lipid metabolism, and body weight in pre-clinical studies. Unlike white fat cells, which store the body's surplus energy as fat, [brown fat](#) cells in BAT can convert some of this energy (lipids and glucose) into large amounts of heat when the body is exposed to cold, thus burning the energy surplus rather than storing it. Estimates suggest that fully activated BAT contributes up to 20% of total [energy expenditure](#). Earlier studies have shown that BAT activity is reduced in obese adults.

South Asians have a much greater risk of developing [type 2 diabetes](#) than do white Caucasians. Moreover, in south Asians, type 2 diabetes develops at a far younger age, and at much lower body mass index (BMI). But, until now, little was known about the causes underlying these differences.

In this study, researchers from the Netherlands compared resting energy expenditure and BAT volume and activity in 12 healthy lean south Asian men (aged about 25 years) and 12 matched white Caucasian men after exposure to cold using 18F-fluorodeoxyglucose positron emission tomography and computed tomography (18F-FDG-PET-CT) scans*. The effect of cold exposure on non-shivering thermogenesis and plasma lipid levels was also measured.

Strikingly, results showed that resting energy expenditure was markedly lower (-32%) in south Asian participants than in Caucasians, as was the volume of metabolically active BAT (-34%). These findings were in line with the significantly higher shiver temperature noted in south Asians (10.9°C vs 8.9°C) upon cold exposure, despite their higher total fat mass. South Asians also had lower cold-induced non-shivering thermogenesis compared with Caucasians.

According to lead authors Mariëtte Boon and Patrick Rensen from Leiden University Medical Center in the Netherlands, "Our findings suggest that making more BAT or increasing its activity could have great therapeutic potential in south Asians, helping them increase the clearance of glucose and fatty acids and converting surplus white fat into heat, potentially lessening the risk of diabetes."**

They add, "It has recently been shown that BAT can be recruited in humans following 10 days of cold intervention. Future studies should be directed towards the efficacy of this strategy, as well as other options, such as medication, to increase BAT activity. These strategies might finally be used to improve the metabolic phenotype in south Asians."**

Writing in a linked Comment, Michael Symonds from The University of Nottingham in the UK points out, "As south Asians are much more susceptible to metabolic disturbances such as [obesity](#) and diabetes, it is important to establish whether these individuals are resistant to established stimulators of BAT thermogenesis such as chronic [cold exposure](#) or dietary stimulants like capsaicin.¹⁰ This group is now the ideal target for new pharmacological interventions that might offer proof-of-principle that enhanced BAT volume or function

can have long-term health benefits against metabolic disease."

More information:

[www.thelancet.com/journals/lan ...](http://www.thelancet.com/journals/lan...)
[\(13\)70156-6/abstract](http://(13)70156-6/abstract)

Provided by Lancet

APA citation: Less 'brown fat' could help explain why a fifth of the world's population is highly susceptible to type 2 diabetes (2013, November 11) retrieved 20 June 2019 from

<https://medicalxpress.com/news/2013-11-brown-fat-world-population-highly.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.