


Racial disparities persist in diabetes care outcomes

2 December 2013

[Full Text \(subscription or payment may be required\)](#)



Copyright © 2013 [HealthDay](#). All rights reserved.

(HealthDay)—Black patients do not receive the same quality of care, and experience poorer intermediate outcomes in the practice setting of a patient-centered medical home (PCMH), according to research published online Nov. 19 in *Diabetes Care*.

In an effort to assess racial differences in processes of care and intermediate outcomes for diabetes, Joseph A. Simonetti, M.D., of the University of Washington in Seattle, and colleagues retrospectively analyzed data for 1,457 adults receiving care within a PCMH-designated academic practice.

The researchers found that, compared with non-Hispanic white patients, more [black patients](#) had a hemoglobin A1C greater than 9 percent, and fewer black patients achieved a hemoglobin A1C less than 7 percent, low-density lipoprotein (LDL) cholesterol less than 100 mg/dL, or [blood pressure](#) less than 140/90 mm Hg (P

"In summary, despite the implementation of a highly innovative PCMH, we observed [racial differences](#) in both processes of care and intermediate outcomes for patients with diabetes," the authors write.

More information: [Abstract](#)

APA citation: Racial disparities persist in diabetes care outcomes (2013, December 2) retrieved 21 October 2021 from <https://medicalxpress.com/news/2013-12-racial-disparities-persist-diabetes-outcomes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.