


Study: CT scans could bolster forensic database to ID unidentified remains

January 14 2014



This cranium image was constructed from CT scans. Credit: Amanda Hale

A study from North Carolina State University finds that data from CT scans can be incorporated into a growing forensic database to help determine the ancestry and sex of unidentified remains. The finding may also have clinical applications for craniofacial surgeons.

"As forensic anthropologists, we can map specific coordinates on a [skull](#) and use software that we developed – called 3D-ID – to compare those three-dimensional coordinates with a database of biological characteristics," says Dr. Ann Ross, a professor of anthropology at NC State and senior author of a paper describing the work. "That

comparison can tell us the ancestry and sex of unidentified remains using only the skull – which is particularly valuable when dealing with incomplete skeletal remains."

However, the size of the 3D-ID database has been limited by the researchers' access to contemporary skulls that have clearly recorded demographic histories.

To develop a more robust database, Ross and her team launched a study to determine whether it was possible to get good skull coordinate data from living people by examining CT scans.

The University of Pennsylvania Museum's Morton Collection provided the NC State researchers with CT scans of 48 skulls. Researchers mapped the coordinates of the actual skulls manually using a digitizer, or electronic stylus. Then they compared the data from the CT scans with the data from the manual mapping of the skulls.

The researchers found that eight bilateral coordinates on the skull – those found on either side of the head – were consistent for both the CT scans and manual mapping.

"This will allow us to significantly expand the 3D-ID [database](#)," Ross says. "And these bilateral coordinates give important clues to [ancestry](#), because they include cheekbones and other facial characteristics."

However, the five midline coordinates the researchers tested showed inconsistencies between the CT scans and manual mapping. Midline coordinates are those found along the center of the skull, such as the bridge of the nose.

"More research is needed to determine what causes these inconsistencies, and whether we'll be able to retrieve accurate midline

data from CT scans," says Amanda Hale, a former master's student at NC State and lead author of the paper.

This research may also help craniofacial surgeons. "An improved understanding of the flaws in how CT scans map skull features could help surgeons more accurately map landmarks for reconstructive surgery," Hale says.

More information: The paper, "A Geometric Morphometric Validation Study of Computed Tomography Extracted Craniofacial Landmarks," is published in the January issue of the *Journal of Craniofacial Surgery*. [journals.lww.com/jcraniofacial ...ion_Study_of.48.aspx](http://journals.lww.com/jcraniofacial...ion_Study_of.48.aspx)

Abstract

This study investigates the variation between craniofacial landmarks extracted from computed tomography (CT) scans and those collected from direct digitization of dry skulls. Thirteen traditional craniofacial landmarks were obtained from each CT scan using the coordinate option in the software Aviso. These coordinates were then compared to the coordinates digitized directly from the dry skulls as two separate samples and individually. Similarities were found between the two coordinate samples with the first principal component representing only 23.97% of the total variation associated with the data acquisition methods and was found to be statistically significant ($p=0.0223$). Differences were more prevalent along midline landmarks. In contrast, the individual specimen comparisons exhibited the largest amount of variation within symmetric landmarks with bilateral landmarks that were more medially located in the CT sample, but no individual specimens were significantly different (e.g. $p\text{-value}=0.9883$) when comparing both data acquisition modalities. Bilateral coordinates were not found to be significantly different for either analysis ($p\text{-value}=0.4165$). The significant differences found for the entire dataset suggest that the combination of CT extracted and

digitized individuals need to be further explored with respect to the reference frames and sample composition. However, the individual specimen comparison results of this study validate the utility of CT extracted landmarks when used for putative identifications in a forensic setting and when clinically applied.

Provided by North Carolina State University

Citation: Study: CT scans could bolster forensic database to ID unidentified remains (2014, January 14) retrieved 20 September 2024 from <https://medicalxpress.com/news/2014-01-ct-scans-bolster-forensic-database.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.