

# Research shows smoking harms your chances of recovering from fractures

March 12 2014

---

Bone healing cells in non-smokers are of a better quality, more active and quicker at dividing than those of smokers, according to new research.

The study, carried out by United Lincolnshire Hospitals NHS Trust (ULHT) and the University of Lincoln, UK, involved gathering 50 fracture patients who volunteered to allow blood from the area of the fracture to be analysed.

The blood was studied in the laboratory at the University of Lincoln to identify the differences in the quantity and quality of stem cells and molecules that are involved in bone regeneration following fracture.

Trauma and Orthopaedic Consultant at Lincoln County Hospital, Professor Mohammad Maqsood, who had previously completed the first stages of research into how smoking affects the [healing](#) of fractures at cellular and [molecular level](#), said: "We set up the research looking at the factors which affect the healing process of fractures in 2007, which involved collecting the blood from consenting patients, both smokers and non-smokers, who have suffered tibial fractures and had been admitted to Lincoln County Hospital for surgery.

"We also set up a 'smoking machine' which simulates smoking 20 cigarettes per day to study the effects of smoking on the blood in a controlled environment."

The team also included Director of Research and Development at ULHT and Consultant Breast Surgeon, Professor Oleg Eremin; Senior Lecturer, Dr Mohamed El-Sheemy, and PhD researcher Andrew Sloan from the University of Lincoln, and staff from the orthopaedic department at Lincoln County Hospital.

Andrew said: "I was particularly interested in how stem cells can be extracted and isolated from human fracture tissue. I felt a real sense of achievement when we were able to do this, as only one group in the world had written about this methodology previously. After growing and harvesting the [stem cells](#) from the tissue, we were able to closely analyse and monitor how cigarette smoking had a detrimental impact on stem cell growth and development, giving rise to abnormal bone repair. We also showed how the molecules that mediate certain cellular processes were being hampered by the toxic effects of the tobacco smoke and its constituent components."

The results revealed that the bone healing cells in non-smokers were of a superior quality, more active and quicker at dividing than those of smokers, which contributes to a faster [healing process](#).

The team presented the findings of this study across Britain and at conferences in Finland, France, Switzerland and the Czech Republic.

The next steps of the project will involve looking at how the adverse effect of [smoking](#) at cellular and molecular level can be reversed.

Provided by University of Lincoln

Citation: Research shows smoking harms your chances of recovering from fractures (2014, March 12) retrieved 16 March 2026 from <https://medicalxpress.com/news/2014-03-chances-recovering-fractures.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.