

Water from improved sources is not consistently safe

6 May 2014

Although water from improved sources (such as piped water and bore holes) is less likely to contain fecal contamination than water from unimproved sources, improved sources in low- and middle-income countries are not consistently safe, according to a study by US and UK researchers, published in this week's *PLOS Medicine*.

These findings are important as WHO and UNICEF track progress towards the Millennium Development Goals water target using the indicator "use of an improved source": this study shows that assuming that "improved" [water sources](#) are safe greatly overestimates the number of people thought to have access to water from a safe source, suggesting that a large proportion of the world's population still use unsafe water.

The authors, led by Robert Bain and Jamie Bartram from The Water Institute at University of North Carolina, reached these conclusions by comprehensively searching the literature to find appropriate studies that investigated [fecal contamination](#) of all types of drinking water in low- and middle-income countries. The authors then investigated the risk factors and settings where fecal contamination of water sources was most common.

319 studies reporting on 96,737 water samples were included in the analysis and the authors found that overall, although the odds (chance) of fecal contamination were considerably lower for "improved" sources than "unimproved" sources, in 38% of 191 studies, over a quarter of samples from improved sources contained fecal contamination. Protected dug wells in particular were rarely free of fecal contamination. Water sources in low-income countries and rural areas were more likely to be contaminated (each with an odds ratio of 2.37).

As included studies rarely reported stored water quality or sanitary risks, these findings may understate the number of people drinking unsafe

water and have profound implications for public health policy.

The authors propose alternative indicators of safe water such as sanitary measures and water quality and say: "Our review provides strong evidence that by equating "improved" with "safe," the number of people with access to a safe water source has been greatly overstated, and suggests that a large number and proportion of the world's population use unsafe [water](#)."

More information: Bain R, Cronk R, Wright J, Yang H, Slaymaker T, et al. (2014) Fecal Contamination of Drinking-Water in Low- and Middle-Income Countries: A Systematic Review and Meta-Analysis. *PLoS Med* 11(5): e1001644. [DOI: 10.1371/journal.pmed.1001644](https://doi.org/10.1371/journal.pmed.1001644)

Provided by Public Library of Science

APA citation: Water from improved sources is not consistently safe (2014, May 6) retrieved 21 October 2020 from <https://medicalxpress.com/news/2014-05-sources-safe.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.