

# Antibody combined with statin results in further reduction of cholesterol levels

13 May 2014

Among patients with high cholesterol receiving moderate- or high-intensity statin therapy, the addition of the human monoclonal antibody evolocumab resulted in additional lowering of low-density lipoprotein cholesterol (LDL-C) levels, according to a study in the May 14 issue of *JAMA*.

Many [patients receiving moderate- or high-intensity statin therapy](#) are unable to achieve recommended LDL-C goals, and consideration of non-statin therapy for additional LDL-C lowering has been recommended. In phase 2 studies, evolocumab use was associated with reduced LDL-C levels in patients receiving statin therapy, according to background information in the article.

Jennifer G. Robinson, M.D., M.P.H., of the University of Iowa, Iowa City, and colleagues conducted a phase 3 study (LAPLACE-2) in which patients were assigned to 1 of 24 treatment groups in two steps. Initially, patients (n = 2,067) were randomly assigned to a daily, moderate-intensity statin ([atorvastatin](#) [10 mg], simvastatin [40 mg], or rosuvastatin [5 mg]); or high-intensity statin (atorvastatin [80 mg], rosuvastatin [40 mg]). After four weeks, while continuing their statin therapy, patients (n = 1,899) were then randomly assigned to compare the effect of evolocumab (140 mg every 2 weeks or 420 mg monthly) with placebo (every 2 weeks or monthly) or ezetimibe (10 mg or placebo daily, atorvastatin patients only; a nonstatin drug used to treat [high cholesterol](#)). The study was conducted at 198 sites in 17 countries.

Compared with placebo, evolocumab provided clinically equivalent percent reductions in LDL-C levels when administered every 2 weeks (66 percent to 75 percent) and monthly (63 percent to 75 percent) when added to moderate- or high-intensity statin [therapy](#). The additional LDL-C lowering seen with evolocumab was greater than that observed with ezetimibe (up to 24 percent reduction).

Evolocumab was well tolerated, with comparable rates of adverse events compared to placebo and ezetimibe during the 12-week treatment period.

"LAPLACE-2 is to our knowledge the first study to demonstrate that the addition of evolocumab results in similar percent reductions in LDL-C and achieved LDL-C levels regardless of stable baseline statin type, dose, or intensity, across 3 commonly prescribed statins and a broad range of doses," the authors write.

**More information:** "Further studies are needed to evaluate longer-term clinical outcomes and safety of this approach for LDL-C lowering." [DOI: 10.1001/jama.2014.4030](#)

Provided by The JAMA Network Journals

APA citation: Antibody combined with statin results in further reduction of cholesterol levels (2014, May 13) retrieved 9 May 2021 from <https://medicalxpress.com/news/2014-05-antibody-combined-statin-results-reduction.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*