


Combo of tools IDs alcohol use in transplant patients

23 May 2014

Copyright © 2014 [HealthDay](#). All rights reserved.



(HealthDay)—The Alcohol Use Disorders Identification Test for alcohol consumption (AUDIT-c) combined with urinary ethyl glucuronide (uEtG) testing improves the detection of alcohol consumption in liver transplant candidates and recipients, according to a study published online April 2 in *Liver Transplantation*.

Salvatore Piano, M.D., from the University of Padova in Italy, and colleagues evaluated uEtG, AUDIT-c, serum ethanol, urinary ethanol, carbohydrate-deficient transferrin (CDT), and other indirect markers of [alcohol consumption](#) in 121 [liver transplant](#) candidates (LTCs) and liver transplant recipients (LTRs).

The researchers found alcohol consumption in 30.6 percent of patients. The strongest marker of alcohol consumption was uEtG (odds ratio, 414.5; P

"In conclusion, the association of AUDIT-c and uEtG improves the detection of alcohol consumption in LTCs and LTRs," the authors write.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

APA citation: Combo of tools IDs alcohol use in transplant patients (2014, May 23) retrieved 4 December 2021 from <https://medicalxpress.com/news/2014-05-combo-tools-ids-alcohol-transplant.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.