

Common treatment of certain autoimmune disease does not appear effective

July 15 2014

Among patients with the systemic autoimmune disease primary Sjögren syndrome, use of hydroxychloroquine, the most frequently prescribed treatment for the disorder, did not improve symptoms during 24 weeks of treatment compared with placebo, according to a study in the July 16 issue of *JAMA*.

Primary Sjögren [syndrome](#) is characterized by mouth and eye dryness, pain, and fatigue, with systemic manifestations occurring in approximately one-third of [patients](#). Despite the wide use of [hydroxychloroquine](#) in clinical practice, evidence regarding its efficacy is limited, according to background information in the article.

Jacques-Eric Gottenberg, M.D., Ph.D., of the Universite de Strasbourg, Strasbourg, France, and colleagues randomly assigned 120 patients with primary Sjögren syndrome to receive hydroxychloroquine or placebo for 24 weeks. All patients (treatment and placebo) were prescribed hydroxychloroquine between weeks 24 and 48. The study was conducted at 15 university hospitals in France.

At 24 weeks, the proportion of patients meeting the primary end point (30 percent or greater reduction in 2 of 3 scores evaluating dryness, pain, and fatigue) was 17.9 percent (10/56) in the hydroxychloroquine group and 17.2 percent (11/64) in the [placebo group](#). Use of hydroxychloroquine was also not associated with improvement in certain lab tests used to monitor the activity of Sjögren (anti-SSA antibodies, IgG levels). During the first 24 weeks, there were 2 serious adverse

events in the hydroxychloroquine group and 3 in the placebo group. In the last 24 weeks, there were 3 in the hydroxychloroquine group and 4 in the placebo group.

"In the present study, hydroxychloroquine did not demonstrate efficacy for the main disabling symptoms—dryness, pain, and fatigue—of primary Sjögren syndrome compared with [placebo](#). [This trial] extends the negative results of the only previous controlled crossover trial, which included 19 patients and confirms that previous open trials might have overestimated the [therapeutic efficacy](#) of hydroxychloroquine in primary Sjögren syndrome," the authors write.

"Further studies are needed to evaluate longer-term outcomes."

More information: [DOI: 10.1001/jama.2014.7682](https://doi.org/10.1001/jama.2014.7682)

Provided by The JAMA Network Journals

Citation: Common treatment of certain autoimmune disease does not appear effective (2014, July 15) retrieved 20 September 2024 from <https://medicalxpress.com/news/2014-07-common-treatment-autoimmune-disease-effective.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--