

Bacteria growing less susceptible to common antiseptic

August 13 2014

Bacteria that cause life-threatening bloodstream infections in critically ill patients may be growing increasingly resistant to a common hospital antiseptic, according to a recent study led by investigators at Johns Hopkins. The study was published in the September issue of *Infection Control and Hospital Epidemiology*, the journal of the Society for Healthcare Epidemiology of America.

Chlorhexidine gluconate (CHG) has been increasingly used in hospitals in light of recent evidence that daily antiseptic baths for patients in intensive care units (ICUs) may prevent infections and stop the spread of healthcare-associated infections. The impact of this expanded use on the effectiveness of the disinfectant is not yet known.

"Hospitals are appropriately using chlorhexidine to reduce infections and control the spread of antibiotic-resistant organisms," said Nuntra Suwantararat, MD, lead author. "However, our findings are a clear signal that we must continue to monitor [bacteria](#) for emerging antiseptic resistance as these antibacterial washes become more widely used in hospitals."

In the study, [investigators](#) compared bacterial resistance between cultures from patients in eight ICUs receiving daily antiseptic washes to patients in 30 non-ICUs who did not bathe daily with CHG. Bacterial cultures obtained from patients with regular antiseptic baths showed reduced susceptibility to CHG when compared with those from patients who did not have antiseptic baths. Regardless of unit protocol, 69

percent of all bacteria showed reduced CHG susceptibility, a trend that requires vigilant monitoring.

"The good news is that most bacteria remain vulnerable to CHG, despite the reduced susceptibility. Daily baths with a CHG solution remain effective against life-threatening bloodstream infections," said Suwantararat.

The investigators caution that the clinical implications of their findings remain unclear. For example, antibiotic susceptibility tests are commonly used to determine whether patients will respond to antibiotic treatment. A similar correlation between antiseptic susceptibility and response to an antiseptic are not as well defined. Identifying particular bacteria and settings in which these bacteria will not respond to [antiseptic](#) agents used in hospitals is an important next step.

More information: Nuntra Suwantararat, Karen C. Carroll, Tsigereda Tekle, Tracy Ross, Lisa L. Maragakis, Sara Cosgrove, Aaron M. Milstone. "High Prevalence of Reduced Chlorhexidine Susceptibility in Organisms Causing Central Line-Associated Bloodstream Infections." *Infection Control and Hospital Epidemiology* [35:9] (September 2014).

Provided by Society for Healthcare Epidemiology of America

Citation: Bacteria growing less susceptible to common antiseptic (2014, August 13) retrieved 25 April 2024 from

<https://medicalxpress.com/news/2014-08-bacteria-susceptible-common-antiseptic.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--