

# Researchers develop the first neurophysiological model of how humans sense wetness

2 October 2014

Human sensitivity to wetness plays a role in many aspects of daily life. Whether feeling humidity, sweat or a damp towel, we often encounter stimuli that feel wet. Though it seems simple, feeling that something is wet is quite a feat because our skin does not have receptors that sense wetness. The concept of wetness, in fact, may be more of a "perceptual illusion" that our brain evokes based on our prior experiences with stimuli that we have learned are wet.

So how would a person know if he has sat on a wet seat or walked through a puddle? Researchers at Loughborough University and Oxygene Research proposed that wetness perception is intertwined with our ability to sense cold temperature and tactile sensations such as pressure and texture. They also observed the role of A-nerve fibers—sensory nerves that carry temperature and tactile information from the [skin](#) to the brain—and the effect of reduced nerve activity on wetness perception. Lastly, they hypothesized that because hairy skin is more sensitive to thermal stimuli, it would be more perceptive to wetness than glabrous skin (e.g., palms of the hands, soles of the feet), which is more sensitive to tactile stimuli.

Davide Filingeri et al. exposed 13 healthy male college students to warm, neutral and cold wet stimuli. They tested sites on the subjects' forearms (hairy skin) and fingertips (glabrous skin). The researchers also performed the wet stimulus test with and without a nerve block. The nerve block was achieved by using an inflatable compression (blood pressure) cuff to attain enough pressure to dampen A-nerve sensitivity.

They found that wet perception increased as temperature decreased, meaning subjects were much more likely to sense cold wet stimuli than warm or neutral wet stimuli. The research team

also found that the subjects were less sensitive to wetness when the A-nerve activity was blocked and that hairy skin is more sensitive to wetness than glabrous skin. These results contribute to the understanding of how humans interpret wetness and present a new model for how the brain processes this sensation.

"Based on a concept of perceptual learning and Bayesian perceptual inference, we developed the first neurophysiological model of cutaneous wetness sensitivity centered on the multisensory integration of cold-sensitive and mechanosensitive skin afferents," the research team wrote. "Our results provide evidence for the existence of a specific information processing model that underpins the neural representation of a typical wet stimulus."

**More information:** "Why wet feels wet? A neurophysiological model of human cutaneous wetness sensitivity." Davide Filingeri , Damien Fournet , Simon Hodder , George Havenith *Journal of Neurophysiology*, Published 15 September 2014 Vol. 112 no. 6, 1457-1469 DOI: 10.1152/jn.00120.2014

Provided by American Physiological Society

APA citation: Researchers develop the first neurophysiological model of how humans sense wetness (2014, October 2) retrieved 20 October 2020 from <https://medicalxpress.com/news/2014-10-neurophysiological-humans.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*