


Obesity fuels silent heart damage

November 20 2014



Heart diagram. Credit: Wikipedia

Using an ultrasensitive blood test to detect the presence of a protein that heralds heart muscle injury, researchers from Johns Hopkins and elsewhere have found that obese people without overt heart disease experience silent cardiac damage that fuels their risk for heart failure down the road.

The findings of the federally funded study, published ahead of print in the *Journal of the American College of Cardiology: Heart Failure*,

challenge the commonly held belief that much of the cardiovascular disease seen in severely overweight people is driven by diabetes and high blood pressure, both well-known cardiac [risk factors](#) and both occurring frequently among the obese.

Specifically, the research showed that obese people had elevated levels of a heart enzyme known as troponin T, released by injured [heart muscle](#) cells. Increases in levels of this enzyme corresponded to increases in people's body mass index (BMI)—a measure of body fat based on a person's weight-to-height ratio. Levels of the enzyme rose proportionally as BMI went up.

Troponin T is the gold standard for diagnosing acute or recent heart attacks and is widely used in emergency rooms to test patients with chest pain and other symptoms suggestive of a heart attack. The test used in the current study works in much the same way, but is calibrated to detect [troponin levels](#) far below the ranges of the clinical test for diagnosing a [heart attack](#).

"Obesity is a well-known 'accomplice' in the development of heart disease, but our findings suggest it may be a solo player that drives [heart failure](#) independently of other risk factors that are often found among those with excess weight," says lead investigator Chiadi Ndumele, M.D., M.H.S., an assistant professor at the Johns Hopkins Ciccarone Center for the Prevention of Heart Disease. "The direct relationship we found between obesity and subclinical heart damage is quite potent and truly concerning from a public health standpoint given the growing number of obese people in the United States and worldwide."

For the study, investigators measured the BMIs and cardiac troponin levels of more than 9,500 heart disease-free men and women, aged 53 to 75, living in Maryland, Mississippi, North Carolina and Minnesota. The researchers then tracked the participants' health for more than 12 years.

During the follow-up, 869 people developed heart failure.

People who were severely obese—those with a BMI above 35—had more than twice the risk of developing heart failure, compared with people of normal weight, the researchers found. That risk rose incrementally with BMI, growing by 32 percent for every five-unit increase in BMI. Thus, a 6-foot, 225-pound man with a BMI of 30 was 32 percent more likely to develop heart failure than a 6-foot, 188-pound man with a BMI of 25. All people with elevated troponin levels, regardless of BMI, had higher risk of developing heart failure over a decade. In other words, extra weight and high troponin each independently signaled higher [heart disease risk](#).

When the researchers calculated the combined effects of elevated troponin and severe obesity, the predictive power was striking. Severely obese people with elevated troponin levels were nine times more likely to develop heart failure than people with normal weight and undetectable troponin levels. The elevated risk persisted even when investigators accounted for other possible causes of heart damage, including diabetes, hypertension and high cholesterol.

Public health experts deem heart failure—a condition in which the heart muscle doesn't pump efficiently—a looming epidemic. The disease has been on a steady rise and is expected to affect one in five adults by 2030.

Ndumele and team say the findings underscore the dangers of obesity and should be heeded as an alarm bell for clinicians to monitor their obese patients rigorously for emerging signs [heart disease](#).

"These results are a wake-up call that obesity may further fuel the growing rate of heart failure, and clinicians who care for [obese people](#) should not be lulled into a false sense of security by the absence of traditional risk factors, such as high cholesterol, diabetes and

hypertension," says Roger Blumenthal, M.D., director of the Johns Hopkins Ciccarone Center for the Prevention of Heart Disease. "Obese people, even when free of cardiovascular symptoms, should be monitored for the earliest signs of heart failure and counseled on ways to improve their lifestyle habits."

The investigators say their next step is to study the precise mechanism by which obesity causes subclinical heart muscle damage, and whether reduction in weight would lower the risk for [heart](#) failure.

Provided by Johns Hopkins University School of Medicine

Citation: Obesity fuels silent heart damage (2014, November 20) retrieved 19 September 2024 from <https://medicalxpress.com/news/2014-11-obesity-fuels-silent-heart.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--