


Dopamine leaves its mark in brain scans

21 November 2014



Left: If the visual cortex of the brain is active, the BOLD signal increases without dopamine. The activities of gamma waves, individual groups of nerve cells (MUA) as well as the blood flow in the area (CBF) also increase. Centre: Under the influence of dopamine, the BOLD signal decreases. The gamma-waves and the activity of the nerve cells, however, remain constant. The blood flow even increases. Right: Active regions (red) in the visual cortex of the brain. Credit: MPI f. Biological Cybernetics / D. Zaldivar

Researchers use functional magnetic resonance imaging (fMRI) to identify which areas of the brain are active during specific tasks. The method reveals areas of the brain, in which energy use and hence oxygen content of the blood changes, thus indirectly showing which cell-populations are particularly active at a given moment. Researchers from the Max Planck Institute for Biological Cybernetics in Tübingen now demonstrate that activity induced by signalling molecules such as dopamine may yield hitherto unpredictable up or down modulations of the fMRI signals, with the result that the neural and vascular responses dissociate. In such cases, far more precise data can be obtained when fMRI is combined with concurrent measurements of cerebral blood flow.

When you work hard, you breathe heavily. The same applies to [nerve cells](#). When neurons fire, they consume more oxygen that is being delivered through blood. To ensure that no deficiency

occurs, an oversupply of oxygenated blood is immediately transported to active regions of the [brain](#). As a result, the [oxygen content](#) of the blood rises in those areas. In a [magnetic resonance](#)-scanner, this process is measured in the form of a blood oxygenation level dependent (BOLD) signal. When the activity of nerve cells increases, the BOLD signal increases too - in theory.

However, external influences such as mood, age, drugs, and food can alter BOLD signals and thereby change the interpretation of fMRI results. Moreover, the results are also affected by different brain states such as attention, memory and reward. "There is no absolute correlation between [neuronal activity](#) and BOLD signals. Consequently, our ability to interpret the signals from fMRI scans is limited," says Daniel Zaldivar of the Max Planck Institute for Biological Cybernetics, describing the motivation starting point of his research. Together with his colleagues, he studied how nerve cells in the visual cortex of macaque monkeys respond to visual stimuli when the brain is simultaneously under the influence of dopamine. The surprising result: although the activity of the nerve cells increases, the BOLD signal decreases by about 50 percent. This can lead the viewer of a brain scan to erroneously conclude that these neurons are less active.

"Dopamine presumably causes active cells to consume more oxygen than can be delivered," says Zaldivar. Paradoxically, dopamine ramps up neurons' activity to such a degree that the BOLD signal shows exactly the opposite of what is really happening. Under the influence of dopamine and probably other neuromodulators, changes in the BOLD signal alone are therefore not sufficient to draw conclusions about the activity of neuronal cells.

Measurements of [cerebral blood flow](#) in combination with BOLD and neurophysiology offer better insight into the changes of energy metabolism and help to draw better conclusion about the neuronal cells activity. That is because

cerebral blood flow provides more direct information about the delivery of oxygen. Interestingly, Zaldivar and colleagues found that under the influence of dopamine, [blood flow](#) increased. These results lead to the conclusion that the increase along with the neural activity is driven by increased energy use.

"If we can improve our understanding of how BOLD signals change under the influence of neuromodulators, we may be able to interpret brain scans better and detect problems at an earlier stage," says Zaldivar. In schizophrenia patients, for example, the dopamine system in the brain is poorly regulated. If scientists knew what impact neuromodulators such as [dopamine](#) have on brain scan images, it might be possible to diagnose such illnesses earlier. "Before drawing conclusions about neuronal activity from BOLD signals, we first need to know what influence neuromodulators have on the images," says Zaldivar.

More information: Daniel Zaldivar, Alexander Rauch, Kevin Whittingstall, Nikos K. Logothetis, Jozien Goense, Dopamine-induced dissociation of BOLD and neural activity in macaque visual cortex, *Current Biology*, 20 November 2014

Provided by Max Planck Society

APA citation: Dopamine leaves its mark in brain scans (2014, November 21) retrieved 30 October 2020 from <https://medicalxpress.com/news/2014-11-dopamine-brain-scans.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.