


Facial motion activates a dedicated network within the brain, research shows

8 January 2015



A network of face processing areas within the macaque brain (red and yellow, shown from the side) responds selectively to face motion over other kinds of motion, researchers discovered. Their work also revealed a new face patch within this network (circled), which they call the middle dorsal patch. Credit: Laboratory of Neural Systems at The Rockefeller University/Current Biology

A face is more than a static collection of features. A shift in gaze, a tightening of the lips, a tilt of the head, these movements convey important clues to someone's state of mind. Scientists know that two particularly social and visual creatures, humans and rhesus macaque monkeys, have a network of small areas within their brains that become active when shown still images of faces. But it hasn't been clear if the same areas are responsible for processing changing expressions and other facial movements.

New research by Clark Fisher, a graduate student in Assistant Professor Winrich Freiwald's Laboratory of Neural Systems at The Rockefeller University confirms that they are. Fisher and Freiwald, building on earlier work by graduate student Pablo Polosecki, found that these face-

processing areas, which they call patches, within a macaque's [brain](#) appear to respond selectively to the motion of [faces](#), while reacting much less to other types of motion.

"Based on input from the eyes, the brain's visual system reconstructs reality, a process that starts out with attributes, such as form, motion and color, that the brain then integrates to build meaningful representations of the world," Freiwald says. "This work reveals how the macaque face processing system reunites two individual streams of visual information - face form and face motion - as it recreates the social reality of a face as a macaque sees it."

Freiwald has been studying brain patches linked to facial recognition for over a decade, and was a contributor to a recent report by the National Research Council that evaluated the science behind eyewitness identification. In previous work with his colleague Doris Tsao, now at the California Institute of Technology, he found that a network of interconnected brain patches found along a deep groove in the sides of the macaque brain form a specialized system for processing faces, just as had been shown in an analogous part of the human brain.

While the system for processing still faces appears similar between humans and macaques, studies of facial motion turned up inconsistent results between the species. Human research said, yes, face patches respond specifically to facial motion, with one patch even reacting more strongly to moving faces than to still ones. However, work with macaques suggested, no, this area of the brain seems to respond to facial motion just because it is motion, not because faces are involved. Given that humans and macaques are close relatives in evolutionary terms, this discrepancy perplexed Freiwald and Fisher.

Brain anatomy makes this puzzle difficult to solve

because the patches of interest are located in a part among the patches, Freiwald says. By better of the macaque brain, the superior temporal sulcus, understanding what they do, individually and that also contains areas that respond to motion in together, we will be able to better understand social general. This creates a challenge for researchers behavior and its many disorders, he says. trying to interpret what is going on in this part of a macaque brain when it sees a moving face.

To tease out the distinction between a response to back, these actions all depend on the brain's ability general motion and one specific to faces, Fisher to represent a face," Freiwald says. "It is possible used high resolution scans, known as functional that the patches' dual sensitivity to form and motion magnetic resonance imaging, to record changes in goes beyond simply integrating those attributes, and hints at how we become aware of and interact blood flow in and around the superior temporal with a fellow being." sulcus.

He showed the macaques a variety of movies of The study is being published in *Current Biology*.

fellow macaques making expressions - Provided by Rockefeller University
aggressively baring their impressive canine teeth,
pleasantly smacking their lips, and making faces as
recognizable to macaques as a smile and a
grimace would be to a human. All the while, the
macaque's head moves, as if turning to address
others around it. The researchers made these
movies themselves by filming macaques in front of
a blue screen. For comparison, Fisher also showed
the macaques images of static faces, as well as
moving and static objects, such as toys. All five of
the previously known facial patches responded
preferentially to moving faces over moving objects,
suggesting that [macaques](#) have neural machinery
dedicated to processing both faces and the
movement associated with them.

The scans also revealed a surprise: A sixth,
previously unknown face patch, which the scientists
refer to as the middle dorsal patch. "The middle
dorsal patch turned up sporadically in response to
static faces, but as soon as we started showing
moving faces, it reliably appeared in each brain
hemisphere for every macaque," Fisher said. "A
similar movement-preferring patch has been
identified in humans, and it appears that the middle
dorsal patch could be the macaque equivalent."

Their experiments also uncovered a new puzzle;
some of the patches were more responsive to
normal movies depicting natural motion over
movies containing the same series of movements
jumbled together randomly. The other patches
"preferred" the unnatural jumbled movies. It's not
clear why, but it is evidence of division of labor

APA citation: Facial motion activates a dedicated network within the brain, research shows (2015, January 8) retrieved 13 June 2021 from <https://medicalxpress.com/news/2015-01-facial-motion-dedicated-network-brain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.