


New insights into underlying cellular mechanisms of information processing in the brain

18 February 2015



A, B, Schematic diagram showing SVs at different priming states located at different distance to the Ca²⁺ source. Only the SVs that locate close to the Ca²⁺ source can be released by AP. We propose that either a single AZ contains a mixture of fast and slow SVs (A) or that there are individual AZs within the calyx that contain only fast or only slow pool SVs (B). In both cases, we propose that distal SVs are rapidly converted to fast pool SVs for AP-evoked release to maintain signaling at high firing rates. Credit: Max Planck Florida Institute for Neuroscience; Dr. Samuel M. Young, Jr.

Researchers at the Max Planck Florida Institute for Neuroscience and the Pasteur Institute have uncovered a key factor in regulating information transmittal during the early stages of auditory processing.

Synapses transmit information from one neuron to another in the form of synaptic vesicles containing neurotransmitters. Continuous release of neurotransmitters is essential to maintain

communication between neurons. To better understand a number of neurological disorders, we need a better understanding of how [synapses](#) continuously relay information between neurons. A new study has discovered a key factor in regulating this continual communication is the proximity of synaptic vesicles next to voltage gated [calcium channels](#) within synapses.

- The human body consists of almost 100 billion neurons that contain synapses, point-to-point contacts for information transfer. Every function, feeling and action depends on the ability of synapses within neurons to maintain continuous communication with each other.
- Synapses transmit information in the form of synaptic vesicles that contain specific chemical messengers called neurotransmitters. The continuous release of neurotransmitters is essential to maintain communication between neurons.
- To better understand and address a number of neurological disorders, we need a better understanding of how synapses can continuously relay information between neurons.
- A new study has discovered that a key factor in regulating this continual communication is the proximity of synaptic vesicles next to voltage gated calcium channels within synapses.

Synaptic vesicles in cell-to-cell communication

While synapses contain hundreds to thousands of synaptic vesicles, when a specific signal is received by the neuron, only a fraction of synaptic vesicles that exist in what is called the readily releasable pool are discharged from one neuron to another. Previous studies have demonstrated that within the

readily releasable pool, synaptic vesicles have different specific kinetic properties of release that impact the type and bandwidth of information that can be relayed in response to stimulation. What was not known was the cellular mechanisms that regulate the release of synaptic vesicles from the readily releasable pool to support the early stages of auditory processing.

New findings

In their February publication in the *Journal of Neuroscience*, the authors of the manuscript report that a dominant factor regulating the properties of [synaptic vesicle](#) release supporting the early stages of auditory processing is the distance between the synaptic vesicles and voltage gated calcium channels within a synapse. The authors characterized the readily releasable pool of synaptic vesicles at the calyx of Held, a critical component of [auditory processing](#), employing several different techniques that allowed them to investigate the pre-synaptic mechanisms of informational transmittal. This study identifies the critical mechanism is the regulation of Ca²⁺ influx through voltage gated calcium channels and release of neurotransmitters by synaptic vesicles. These findings are important for understanding the mechanisms of synaptic transmission, specifically for neuronal circuits that rely on fast, continuous synaptic transmission.

"It is becoming apparent that the underlying cause of most neuropsychiatric or neurodegenerative diseases is a dysfunction of the synapse," explained Dr. Young. "Identifying what factors are involved in proper synaptic transmission and neural circuit function have tremendous potential as therapies for neurological disorders or brain injury."

According to Dr. Young, the future goals of this project are to uncover molecular mechanisms that allow synapses to sustain [synaptic transmission](#) over a wide range of activity levels to allow for proper information processing by the neuronal circuit in which they are embedded.

Provided by Max Planck Florida Institute for Neuroscience

APA citation: New insights into underlying cellular mechanisms of information processing in the brain (2015, February 18) retrieved 15 October 2019 from <https://medicalxpress.com/news/2015-02-insights-underlying-cellular-mechanisms-brain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.