

New biomarker identified in women with mental illness

19 June 2015



Credit: George Hodan/Public Domain

Psychiatric disorders can be difficult to diagnose because clinicians must rely upon interpreted clues, such as a patient's behaviors and feelings. For the first time, researchers at University of California, San Diego School of Medicine report identifying a biological marker: the over-production of specific genes that could be a diagnostic indicator of mental illness in female psychiatric patients.

The study was published this week in the journal *EBioMedicine*.

Researchers found that the gene XIST, which is responsible for inactivating one of the two copies of the X chromosome in cells that store genetic material, works overtime in [female patients](#) with [mental illnesses](#), such as bipolar disorder, major depression and schizophrenia.

The study suggests that over-production of XIST and genes from the inactive X chromosome are common denominators in the development of psychiatric disorders in patients with rare chromosome disorders, such as Klinefelter

syndrome and Triple X syndrome, and in the general population of female psychiatric patients.

"There has been an utmost urgency to identify biomarkers for mental illness that could significantly impact research and drug development," said Xianjin Zhou, PhD, assistant professor in the Department of Psychiatry at UC San Diego School of Medicine and lead author.

The study was conducted on 60 lymphoblastoid cell lines from female patients, most of whom had a family history of mental illness. Approximately 50 percent of the female patients exhibited abnormally higher levels of XIST and other genes related to the X chromosome.

Zhou and his team said reversing the abnormal activity of the inactive X chromosome in [patients](#) suffering from mental illness may offer a potential new strategy for treating [psychiatric disorders](#).

"Our results indicate that a large subpopulation of female [psychiatric patients](#) from the [general population](#) may have abnormal function of the inactive X chromosome," said Zhou. "These results are powerful in that early diagnosis of mental illness could possibly happen with a simple blood test, leading to better interventions, therapy and treatment options."

Provided by University of California - San Diego

APA citation: New biomarker identified in women with mental illness (2015, June 19) retrieved 16 October 2021 from <https://medicalxpress.com/news/2015-06-biomarker-women-mental-illness.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.