


Neural basis of multitasking identified

September 1 2015



Plots that show connection strength between nodes in the brain during the control

task (top) and the memory task (bottom). Brain regions are arrayed on both the X and Y axes; the strength of their connection is indicated by the color of the coordinate where they meet. Credit: *Proceedings of the National Academy of Sciences*

What makes someone better at switching between different tasks? Looking for the mechanisms behind cognitive flexibility, researchers at the University of Pennsylvania and Germany's Central Institute of Mental Health in Mannheim and Charité University Medicine Berlin have used brain scans to shed new light on this question.

By studying networks of activity in the brain's [frontal cortex](#), a region associated with control over thoughts and actions, the researchers have shown that the degree to which these networks reconfigure themselves while switching from task to task predicts people's [cognitive flexibility](#).

Experiment participants who performed best while alternating between a memory test and a control test showed the most rearrangement of connections within their frontal cortices as well as the most new connections with other areas of their brains.

A more fundamental understanding of how the brain manages multitasking could lead to better interventions for medical conditions associated with reduced [executive function](#), such as autism, schizophrenia or dementia.

Danielle Bassett, the Skirkanich Assistant Professor of Innovation in Penn's School of Engineering and Applied Science, is senior author on the study. Mannheim's Urs Braun and Axel Schäfer were the lead authors. The research also featured work from Andreas Meyer-Lindenberg and Heike Tost of Mannheim, Henrik Walter of Charité, and others.

It was published in the *Proceedings of the National Academy of Sciences*.

Rather than looking at the role a single region in the brain plays, Bassett and colleagues study the interconnections between the regions as indicated by synchronized activity. Using fMRI, they can measure which parts of the brain are "talking" to one another as study participants perform various tasks. Mapping the way this activity network reconfigures itself provides a more holistic view of how the brain operates.

"We try to understand how dynamic flexibility of brain networks can predict cognitive flexibility, or the ability to switch from task to task," Bassett said. "Rather than being driven by the activity of single brain areas, we believe executive function is a network-level process."

A previous study that Bassett led showed that people who could more quickly "disconnect" their frontal cortices did better on a task that involved pressing keys that corresponded to color-coded notes on a screen. The high level decision-making associated with the frontal cortex's cognitive control wasn't as critical to playing the short sequences of notes, so those who still engaged this part of the brain were essentially overthinking a simple problem.

In the new experiment, lead by Andreas Meyer-Lindenberg of Mannheim, 344 participants alternated between a working [memory task](#) designed to engage the frontal cortex and a control task. The easy task involved pressing the corresponding button as a sequence of numbers appeared on a screen one by one. The hard task also involved a sequence of numbers on a screen, but participants had to press the button that corresponded to the number that appeared two places back in the series each time they saw a new one.

Urs Braun and Axel Schäfer, the lead authors on the paper, collaborated

with Bassett, who has developed novel tools from network science to distill evolving brain connections. They used these tools to map how participants' [brain](#) activities rearranged during each block of the working memory task, each block of the control task and blocks in between where participants switched gears.

"The nodes in the network that are most involved in reconfigurations are cognitive control areas in the frontal cortex," Bassett said. "More flexibility within the frontal cortex meant more accuracy on the memory task, and more consistent connectivity between the frontal cortex and other regions was even more predictive."

While the predictive strength of this reconfiguration suggests that it is only one of several processes involved in successful task switching, it plays a core role.

"It doesn't account for a huge amount variance," Bassett said, "but it suggests that this kind of reconfiguration is a fundamental aspect of cognitive flexibility."

More information: Dynamic reconfiguration of frontal brain networks during executive cognition in humans, Urs Braun, [DOI: 10.1073/pnas.1422487112](#)

Provided by University of Pennsylvania

Citation: Neural basis of multitasking identified (2015, September 1) retrieved 21 September 2024 from <https://medicalxpress.com/news/2015-09-neural-basis-multitasking.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is

provided for information purposes only.