

# Continuation of antibiotics for UTI often inappropriate

17 November 2015

Copyright © 2015 [HealthDay](#). All rights reserved.



(HealthDay)—For patients with urinary tract infections (UTIs), initiation of antibiotics in the emergency department is frequently inappropriate, as is continuation of antibiotics after admission, according to a study published online Nov. 12 in the *Journal of Hospital Medicine*.

Dmitry Kiyatkin, M.D., from the Johns Hopkins Bayview Medical Center in Baltimore, and colleagues reviewed the medical records for all patients admitted to the hospital who initiated treatment for a UTI in the [emergency department](#) during a four-week period.

The researchers found that antibiotic initiation was inappropriate for 59 percent of 94 patients. Continuation of antibiotics after admission was inappropriate for 68 percent of 80 patients.

"In conclusion, we found a high rate of inappropriate antibiotic administration for UTIs that began in the emergency department and continued after admission," the authors write. "Specific guidelines should be developed and validated to direct diagnosis and treatment of UTIs in the emergency department and hospital."

**More information:** [Abstract](#)  
[Full Text \(subscription or payment may be required\)](#)

APA citation: Continuation of antibiotics for UTI often inappropriate (2015, November 17) retrieved 29 November 2021 from <https://medicalxpress.com/news/2015-11-antibiotics-uti-inappropriate.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*