

# Lixisenatide doesn't affect cardiovascular risk in T2DM

3 December 2015



The study was funded by Sanofi, the manufacturer of lixisenatide.

**More information:** [Full Text \(subscription or payment may be required\)](#)

Copyright © 2015 [HealthDay](#). All rights reserved.

(HealthDay)—The addition of lixisenatide to usual care does not impact the rate of major cardiovascular events or other serious adverse events among patients with type 2 diabetes and a recent acute coronary syndrome, according to a study published in the Dec. 3 issue of the *New England Journal of Medicine*.

Marc A. Pfeffer, M.D., Ph.D., from Brigham and Women's Hospital in Boston, and colleagues randomized patients with type 2 diabetes who had had [myocardial infarction](#) or who had been hospitalized for [unstable angina](#) within the prior 180 days to lixisenatide or placebo. A total of 6,068 patients were randomized and followed for a median of 25 months.

The researchers found that a primary end point event (cardiovascular death, myocardial infarction, stroke, or hospitalization for unstable angina) occurred in 13.4 and 13.2 percent of the lixisenatide and placebo groups, respectively (hazard ratio, 1.02; 95 percent confidence interval, 0.89 to 1.17), demonstrating noninferiority (P

"The neutral cardiovascular profile associated with lixisenatide will inform physicians' and patients' decisions regarding the use of this agent as an adjunctive therapy to control the glycosylated hemoglobin level safely," the authors write.

APA citation: Lixisenatide doesn't affect cardiovascular risk in T2DM (2015, December 3) retrieved 6 May 2021 from <https://medicalxpress.com/news/2015-12-lixisenatide-doesnt-affect-cardiovascular-t2dm.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*