

# Hepatitis C tied to increased risk of Parkinson's

December 24 2015

---



Immunohistochemistry for alpha-synuclein showing positive staining (brown) of an intraneural Lewy-body in the Substantia nigra in Parkinson's disease. Credit: Wikipedia

The hepatitis C virus may be associated with an increased risk of

developing Parkinson's disease, according to a study published in the December 23, 2015, online issue of *Neurology*, the medical journal of the American Academy of Neurology. Parkinson's disease is considered the second most common degenerative brain disorder after Alzheimer's disease. Hepatitis C is a liver infection caused by a virus.

"Many factors clearly play a role in the development of Parkinson's disease, including environmental factors," said study author Chia-Hung Kao, MD, China Medical University in Taichung, Taiwan. "This nationwide study, using the National Health Insurance Research Database of Taiwan, suggests that hepatitis caused specifically by the hepatitis C virus may increase the risk of developing the disease. More research is needed to investigate this link."

The World Health Organization estimates that 130 to 150 million people have hepatitis C worldwide. While hepatitis C can lead to serious illness, many people have few symptoms and do not realize they have the virus, especially at first.

The virus is transmitted through sharing needles, needle stick injuries in health care providers and passed on at birth from infected mothers. In Taiwan during the time of the study, blood transfusions were the most common cause of the virus. In the United States, all donated blood has been screened for the virus since 1992.

The study involved 49,967 people with hepatitis and 199,868 people without hepatitis. Participants with hepatitis were placed into three groups: those infected with the hepatitis B virus (71 percent), those with hepatitis C (21 percent), and those who had both viruses (8 percent).

The participants were followed for an average of 12 years to see who developed Parkinson's disease. Of those with hepatitis, 270 developed Parkinson's disease, including 120 people with hepatitis C. Among those

who did not have hepatitis, 1,060 developed Parkinson's disease.

Once researchers controlled for factors such as age, sex, diabetes and cirrhosis, they found that people with hepatitis C were nearly 30 percent more likely to develop Parkinson's disease than the people who did not have hepatitis.

People with hepatitis B and those with both viruses were not more or less likely to develop Parkinson's than those who did not have [hepatitis](#).

Provided by American Academy of Neurology

Citation: Hepatitis C tied to increased risk of Parkinson's (2015, December 24) retrieved 24 April 2024 from <https://medicalxpress.com/news/2015-12-hepatitis-tied-parkinson.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--