

Neurobiology scientists discover new mechanism for detecting sour taste

January 5 2016, by Robert Perkins



New research provides surprising clues as to why some foods taste more sour.

Researchers have discovered a new way that taste cells detect sourness, a sensation that is linked to acidity.

Of the five basic tastes, sour remains the most mysterious—to the point

that the tongue's sour receptor still has not been identified, and the ways in which it receives sour stimuli are still the object of intense research.

"Before 2006, we didn't even know definitively which cells detect sour taste," said Emily Liman, professor of neurobiology at USC Dornsife.


In 2010, Liman identified one of the molecular pathways by which acidic substances trigger the sour taste sensation. Hydrogen ions released by acids directly enter taste cells on your tongue and trigger an electrical signal to the brain.

More than pH

If that were the only way that sour taste was produced, then the lower the pH, the more sour something should taste. However, some substances—such as vinegar—taste more sour than one would expect just based on pH.

A substance's pH is just part of the answer—the concentration of weak acids (such as acetic acid found in vinegar) plays a significant role in how sour it tastes, according to a study published Dec. 1 in *Proceedings of the National Academy of Sciences* by a group of researchers including Liman.

HOW TASTE BUDS WORK



Credit: USC Graphic courtesy of Jonas Töle and David Medzerian.

Previous studies showed that the concentration of a weak acid, in addition to its pH, determines its sourness.

By marking sour taste cells with a fluorescent protein, Liman discovered that there's another way to detect sour tastes. The group identified a potassium channel that makes sour [taste cells](#) extremely sensitive to changes in the cell's intracellular pH.

Making the cell more acidic, too

Like strong acids, weak acids release [hydrogen ions](#) that pass through proteins in the cell membrane, triggering a sour sensation. However, weak acids can go even further, crossing the [cell membrane](#) on their own to actually acidify the fluid inside the cell.

"Taste represents a fairly direct link between sensation and perception, so it's an interesting window into how our body and mind are connected," Liman said. "The mechanism that we discovered not only explains why weak acids taste more [sour](#) than strong acid, but it may also explain how we can detect relatively low concentrations of protons because it predicts that there would an amplification of the initial signal."

More information: Wenlei Ye et al. The K channel K 2.1 functions in tandem with proton influx to mediate sour taste transduction , *Proceedings of the National Academy of Sciences* (2015). [DOI: 10.1073/pnas.1514282112](#)

Provided by University of Southern California

Citation: Neurobiology scientists discover new mechanism for detecting sour taste (2016, January 5) retrieved 19 September 2024 from <https://medicalxpress.com/news/2016-01-neurobiology-scientists-mechanism-sour.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--