


# **Anti-inflammatory drug may prevent rapid aging in people with HIV**

February 22 2016

---



Shokrollah Elahi is lead author of a study that show an anti-inflammatory drug may prevent accelerated aging in HIV-infected individuals.

New research from the University of Alberta's School of Dentistry shows that a commonly used cholesterol-lowering drug could help people with HIV live longer and enjoy an improved quality of life.

The study, Atorvastatin restricts HIV replication in CD4+ T cells by upregulation of p21, published in AIDS (the official journal of the International AIDS Society), shows that the anti-inflammatory drug atorvastatin (Lipitor) may prevent rapid aging associated with the disease.

"HIV-infected individuals experience accelerated aging in different organs as the results of [chronic inflammation](#) and elevated immune activation in them," says Shokrollah Elahi, lead author of the study and immunologist at the School of Dentistry. "If you reduce chronic inflammation, you can enhance their quality of life and expand their life expectancy."

With the availability of [antiretroviral therapy](#), HIV infection has become a chronic disease. But this chronic inflammation state has been linked to rapid aging. As a result, patients now face a new set of challenges such as increased rates of cardiovascular disease, stroke, kidney and liver disease, osteoporosis, neurological disease and non-AIDS-defining malignancies, many of which are more typical of an aging population.

It has been shown that HIV-infected individuals have much higher levels of inflammation within their body than uninfected people do, despite the antiretroviral therapy.

"Human HIV infection is suppressed but not eliminated by the current medication. As a result, viral persistence in the presence of antiretroviral therapy is a major source of inflammation and substantial immune activation, both of which are linked to 'inflammaging'—a concept that contributes to the aging process in HIV-infected individuals," says Elahi. "Our study shows that cholesterol-lowering medication can reduce inflammation and makes CD4 T cells (HIV target cells) less permissible to HIV infection."

The study shows that atorvastatin—the most commonly used cholesterol-lowering medication—effectively reduces [immune activation](#), thereby diminishing the [systemic inflammation](#) that leads to premature or accelerated aging.

Because HIV infection also causes lipid abnormalities such as high cholesterol levels in patients, HIV-infected individuals could benefit from this property of atorvastatin.

The researchers also discovered the drug reduces HIV replication by a novel mechanism, making CD4 T cells more resistant to HIV infection.

Elahi says further clinical trial studies may lead researchers to new therapeutic options to more effectively reduce the inflammation and benefit the health of HIV-infected individuals by preventing accelerated aging.

**More information:** Shokrollah Elahi et al. Atorvastatin restricts HIV replication in CD4+ T cells by upregulation of p21, *AIDS* (2016). [DOI: 10.1097/QAD.0000000000000917](https://doi.org/10.1097/QAD.0000000000000917)

Provided by University of Alberta

Citation: Anti-inflammatory drug may prevent rapid aging in people with HIV (2016, February 22) retrieved 18 April 2024 from <https://medicalxpress.com/news/2016-02-anti-inflammatory-drug-rapid-aging-people.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.