

Antidepressant use begins years before Alzheimer's diagnosis

4 March 2016

Provided by University of Eastern Finland

Antidepressants are frequently prescribed to persons with Alzheimer's disease before the diagnosis, according to a recent study from the University of Eastern Finland. Among persons with Alzheimer's disease, the initiation of antidepressant use was most common during the six months after the Alzheimer's diagnosis, and more frequent than among comparison persons without Alzheimer's disease even 4 years after the diagnosis. The results were published in *International Journal of Geriatric Psychiatry*.

Antidepressant initiation was more frequent among persons with Alzheimer's disease nine years before the diagnosis than among people not diagnosed with Alzheimer's disease. This finding was somewhat surprising. It may be related to more frequent treatment of depressive symptoms because depression has been associated with an increased risk of Alzheimer's disease in previous studies.

Antidepressant use was investigated in the Finnish nationwide MEDALZ study from nine years before until four years after the [diagnosis](#) of Alzheimer's disease. The difference in antidepressant initiations persisted during the entire follow-up period. During the 13-year follow-up period, 42 percent of persons with Alzheimer's disease and 22 percent of persons not diagnosed with the disease initiated antidepressant use. The most commonly used antidepressant group was selective serotonin reuptake inhibitor (SSRI) drugs, followed by mirtazapine.

The register-based study was based on 62,104 Finnish persons with Alzheimer's disease diagnosed between 2005 and 2011.

More information: Arto Puranen et al. Incidence of antidepressant use in community-dwelling persons with and without Alzheimer's disease: 13-year follow-up, *International Journal of Geriatric Psychiatry* (2016). [DOI: 10.1002/gps.4450](https://doi.org/10.1002/gps.4450)

APA citation: Antidepressant use begins years before Alzheimer's diagnosis (2016, March 4) retrieved 19 January 2022 from <https://medicalxpress.com/news/2016-03-antidepressant-years-alzheimer-diagnosis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.