

# Blacks face a higher risk of kidney failure than whites, regardless of genetics

10 March 2016

New research investigates the ties between certain genetic variants and kidney disease in African Americans. The findings, which appear in an upcoming issue of the *Journal of the American Society of Nephrology (JASN)*, suggest that widespread screening for these variants in the black general population is not yet justified.

African Americans have an elevated risk for [chronic kidney disease](#) and [kidney failure](#) compared with European Americans. Studies have shown that much of this risk is due to genetic variations in a gene called apolipoprotein L1 (APOL1), which creates a protein that is a component of HDL, or good cholesterol. These variants arose tens of thousands of years ago in sub-Saharan Africa, and so are present in individuals who have recent sub-Saharan African ancestry. Approximately 5 million African Americans carry APOL1 risk variants; however, not all persons with such variants develop kidney disease.

To get a better sense of how APOL1 genetic risk variants affect [kidney disease](#) and other aspects of health over the long term, Morgan Grams, MD PhD (Johns Hopkins University) and her colleagues evaluated the prognosis and APOL1 status of participants in the Atherosclerosis Risk in Communities (ARIC) study. Among 15,140 ARIC participants followed from 1987-1989 to 2011-2013, 75.3% were white, 21.5% were black/APOL1 low-risk, and 3.2% were black/APOL1 high-risk. "Our study is a population-based cohort following participants over 25 years and thus well suited at assessing a fairly comprehensive set of outcomes among people with the high-risk genotype," said Dr. Grams.

In analyses that adjusted for differences in demographics, blacks had a higher risk for all assessed adverse health events: acute kidney injury, kidney failure, hypertension, diabetes, cardiovascular disease, hospitalization, and death;

however, in analyses that also adjusted for comorbid conditions and socioeconomic status, blacks had a higher risk for hypertension, diabetes, and kidney failure only. When considering only blacks, the APOL1 high-risk variants were linked with a higher risk of kidney failure, but there was high variability in kidney function decline among those with and without the variants.

"We found great variability in [kidney](#) function trajectory, such that most African Americans with the high-risk genotype experienced similar decline as African Americans with the low-risk genotype," said Dr. Grams. "We did find pervasive racial disparities in adverse health outcomes not explained by the APOL1 risk variants, which suggests that interventions to improve health and [health](#) outcomes in African Americans are needed."

**More information:** "Race, APOL1 Risk, and eGFR Decline in the General Population," [DOI: 10.1681/ASN.2015070763](https://doi.org/10.1681/ASN.2015070763)

Provided by American Society of Nephrology

APA citation: Blacks face a higher risk of kidney failure than whites, regardless of genetics (2016, March 10) retrieved 21 September 2019 from <https://medicalxpress.com/news/2016-03-blacks-higher-kidney-failure-whites.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*