


Electrical brain stimulation could support stroke recovery

March 16 2016



Increased fMRI activity and gray matter volume in stroke patients receiving motor rehabilitation plus anodal tDCS compared to motor rehabilitation plus sham treatment. Images are in radiological convention (R, right), and all lesions appear on the left of the image; yellow flash indicates the target for anodal tDCS. (A) Brain regions showing increases in fMRI activity during affected hand movement from baseline to immediately after intervention for the anodal tDCS group versus the sham treatment group. (B) Regions showing greater increases in movement-related fMRI activity for anodal tDCS versus sham group from baseline to 1-month follow-up. (C) Brain regions showing increases in gray matter density from baseline to immediately after intervention for the anodal tDCS group versus the sham treatment group. Credit: C. Allman et al., Science Translational Medicine (2016)

Applying an electric current to the brain can help recovery from stroke, Oxford University researchers have found.

A team from Oxford's Nuffield Department of Clinical Neurosciences, led by Professor Heidi Johansen-Berg and Dr Charlotte Stagg, studied the use of transcranial direct current stimulation (tDCS) to support rehabilitation [training](#). The technique involves placing electrodes on the scalp to pass a constant low current through a particular area of the brain.

In this case, the team used a variant called ipsilesional anodal tDCS, where a positive (anodal) current is applied on the side of the brain where damage has occurred. Anodal stimulation has previously been shown to increase the learning of [motor skills](#) in healthy people. The hope was that this effect could also be demonstrated in stroke patients, using tDCS to reinforce training that helps patients relearn how to use their body.

Professor Heidi Johansen-Berg said: 'For [stroke patients](#), longer and more intensive training leads to greater recovery. However, cost and staff availability limit what can be provided. That means that there is increasing interest in therapies that can be used to boost the effects of training.'

The study included twenty-four volunteers who had had a stroke affecting their hand and arm function, split into two groups. Both groups were given nine days of motor training. One group had tDCS during the training sessions, while the other group acted as a control: they were fitted with electrodes but did not receive tDCS.

Before, and at various times up to three months after the training, the volunteers' motor skills were assessed using established clinical measures to see how much they had improved.

Professor Johansen-Berg said: 'The assessments before the training were used to establish a baseline score for motor skills. Further assessments could then be used to determine what improvement there was above that baseline.

'Three months after training, the group that had received tDCS had improved more on our clinical measures than those in the control group. This showed that the patients who had received tDCS were better able to use their hands and arms for movements such as lifting, reaching and grasping objects.'

MRI scanning also showed that those who had had tDCS had more activity in the relevant brain areas for motor skills than the control group.

Study volunteer Jan said: 'The training was exhausting - like being in the gym every day, but it was huge fun. Even after the first session I felt as

if I could do more, even though I was knackered. That made me go back every day, and I found it easier and easier. [The stimulation] didn't hurt - more like a mild tingle or a static electric shock right on the top of my head. The worst part was that my head itched afterwards!'

She added: 'I have definitely improved and benefited. People who haven't seen me say 'wow - you can move better now'. It definitely helped. I'm just sorry I can't continue with it. It was so nice to meet a team who had such positive attitudes and who told me it was not too late to improve.'

The research team conclude that there is positive evidence for the use of tDCS to aid stroke recovery but caution that the technique must be proved to have long term benefits not only in clinical measurements but also in the ability to carry out tasks important to daily life. Larger studies, they say, will be needed before this approach could enter routine clinical care.

More information: "Ipsilesional anodal tDCS enhances the functional benefits of rehabilitation in patients after stroke," [DOI: 10.1126/scitranslmed.aad5651](https://doi.org/10.1126/scitranslmed.aad5651)

Provided by University of Oxford

Citation: Electrical brain stimulation could support stroke recovery (2016, March 16) retrieved 26 April 2024 from <https://medicalxpress.com/news/2016-03-electrical-brain-recovery.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.