


# Abaloparatide may help prevent fractures in postmenopausal women with osteoporosis

April 3 2016

---



On the left is normal bone and on the right is osteoporotic bone. Credit: International Osteoporosis Foundation

The investigational drug abaloparatide-SC (subcutaneous) may help increase bone mineral density in postmenopausal women and reduce their risk of fracture, new industry-sponsored research suggests. The results of the subgroup analysis within the ACTIVE clinical trial will be presented Friday, April 1, at ENDO 2016, the annual meeting of the Endocrine Society in Boston.

"Abaloparatide-SC increased bone mineral density and reduced the risk

of vertebral and nonvertebral fractures consistently in [postmenopausal women](#) with osteoporosis regardless of their baseline patient characteristics, including age, bone mineral density, and whether or not they had prior fractures," said lead study author Felicia Cosman, MD, endocrinologist and osteoporosis specialist at Helen Hayes Hospital in West Haverstraw, New York, and professor of clinical medicine at Columbia University in New York City.

"If approved by the Food and Drug Administration (FDA), abaloparatide-SC may have the potential to reduce the risk of fractures in postmenopausal women with osteoporosis across a broad range of patient characteristics," said Cosman, who is also a consultant to Radius Health, Inc., in Waltham, Massachusetts.

The researchers investigated patients enrolled in the randomized, double-blind, multinational phase 3 ACTIVE trial to evaluate the efficacy and safety of 80 micrograms of abaloparatide-SC in preventing fractures in otherwise healthy, ambulatory, postmenopausal women with osteoporosis. Overall, 2,463 patient between 49 and 86 years of age were randomized to take one of three medications for 18 months: 80 micrograms of abaloparatide-SC, 20 micrograms of subcutaneous teriparatide (an FDA-approved prescription drug known to increase bone density and strength), or placebo.

At 18 months, abaloparatide-SC significantly increased bone mineral density from baseline at the lumbar spine by 9.2 percent, the total hip by 3.4 percent and the femoral neck by 2.9 percent, compared with placebo. Abaloparatide-SC also reduced new vertebral fractures by 86 percent, nonvertebral fractures by 43 percent and major osteoporotic fractures by 70 percent, compared with placebo. The drug also reduced major osteoporotic fractures by 55 percent, compared with teriparatide.

The authors found that the risk of new vertebral or nonvertebral

fractures dropped consistently among the participants taking abaloparatide-SC, irrespective of their baseline risk factors; and they found no interactions between any of the baseline risk factors and the magnitude of increased bone mineral density buildup by abaloparatide.

Provided by The Endocrine Society

Citation: Abaloparatide may help prevent fractures in postmenopausal women with osteoporosis (2016, April 3) retrieved 20 September 2024 from

<https://medicalxpress.com/news/2016-04-abaloparatide-fractures-postmenopausal-women-osteoporosis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.