

Can your fitness tracker save your life in the ER?

April 5 2016



Heartbeat on an activity tracker/smartphone app. Credit: Alfred Sacchetti, MD

Emergency physicians used a patient's personal activity tracker and

smartphone to identify the time his heart arrhythmia started, which allowed them to treat his new-onset atrial fibrillation with electrical cardioversion and discharge him home. The first case report using information in an activity tracker/smartphone system to assist in medical decision-making was reported online Friday in *Annals of Emergency Medicine*.

"Using the patient's activity tracker - in this case, a Fitbit - we were able to pinpoint exactly when the patient's normal [heart rate](#) of 70 jumped up to 190," said corresponding study author Alfred Sacchetti, MD, FACEP of Our Lady of Lourdes Medical Center in Camden, N.J. "The device told us that the patient's atrial fibrillation was present for only a few hours. That was well within the 48-hour window needed to consider him for rhythm conversion, so we cardioverted him and sent him home."

A 42-year-old patient with a history of seizures but no history of cardiac disease or prior episodes of atrial fibrillation came to the [emergency department](#) following a seizure. He had an irregular heart rate ranging between 130 and 190 beats per minute. He was medicated with oxcarbazepine and diltiazem. His heart rate returned to normal (between 80 and 100 beats per minute) but the atrial fibrillation remained.

The treatment of recent onset atrial fibrillation is electrical cardioversion in any patient who can reliably relate an arrhythmia onset time of within the previous 48 hours. Because the patient was asymptomatic during his current episode of atrial fibrillation, it was not possible to assign an onset time for his arrhythmia.

Emergency department staff accessed the smartphone application connected with his activity tracker and discovered the onset time for his atrial fibrillation was 3 hours prior to coming to the emergency department. After cardioverting the sedated patient, emergency department staff interrogated the smartphone app again, which

accurately recorded the change in heart rate consistent with a rhythm change from [atrial fibrillation](#) to normal rhythm. The patient was discharged home with instructions to follow up with outpatient cardiology.

"Not all activity trackers measure heart rates, but this is the function of most value to medical providers," said Dr. Sacchetti. "Dizziness with a heart rate of 180 would be approached very differently from the same complaint with a heart rate of 30. At present, activity trackers are not considered approved medical devices and use of their information to make medical decisions is at the clinician's own discretion. However, the increased use of these devices has the potential to provide [emergency physicians](#) with objective clinical information prior to the patient's arrival at the emergency department."

More information: "Interrogation of Patient Activity Tracker to Assist Arrhythmia Management", www.annemergmed.com/article/S01600143-8/fulltext

Provided by American College of Emergency Physicians

Citation: Can your fitness tracker save your life in the ER? (2016, April 5) retrieved 26 April 2024 from <https://medicalxpress.com/news/2016-04-tracker-life-er.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.