


Decade of hormone therapy reduces breast cancer recurrence: study

5 June 2016



Micrograph showing a lymph node invaded by ductal breast carcinoma, with extension of the tumour beyond the lymph node. Credit: Nephron/Wikipedia

Doubling the duration of hormone therapy for women with early-stage breast cancer from five to 10 years reduces the risk of recurrence and developing tumors in the other breast, a study showed Sunday.

The study focused on a type of early-stage [breast cancer](#) known as hormone receptor-positive and was presented at the annual meeting of the American Society of Clinical Oncology (ASCO) in Chicago this weekend. It also appears in the *New England Journal of Medicine*.

Postmenopausal women who took the drug letrozole for a decade instead of an often prescribed five-year period saw a 34 percent lower risk of [recurrence](#) or of a new cancer in the other breast than those who received a placebo, according to researchers.

Letrozole is an aromatase inhibitor, which stops

estrogen production in postmenopausal women.

The data suggests that "longer durations of widely-available therapy reduce the risk of cancer recurrence, and prevent second cancers from arising," said Harold Burstein, a breast cancer expert and associate professor of medicine at Harvard Medical School not linked to the study.

Lead study author Paul Goss, director of [breast cancer research](#) at Boston's Massachusetts General Hospital, said women with early stage hormone-receptor positive breast cancer face an "indefinite risk of relapse."

The study involved 1,918 [postmenopausal women](#) in Canada and the United States who had received five years of any one of three [aromatase inhibitor](#) therapies either as initial treatment or after taking tamoxifen, another anti-estrogen drug dating back to the 1970s.

Half of the participants received letrozole after a random selection process, while the other half received a placebo.

After a follow-up period of a little more than six years, just seven percent of participants in the letrozole group saw a recurrence of cancer or a tumor in the other breast, compared to 10.2 percent in the [control group](#).

But only 1.4 percent of the women on letrozole developed a cancer in the breast that had been spared, compared to 3.2 percent of those on the placebo.

As concerns recurrence of the initial cancer—most frequently in the bones, liver or other organs—the difference between the two groups was smaller—5.7 percent in the letrozole group versus 7.1 percent in the control group.

At five years of follow-up, 95 percent of the women

receiving letrozole were free of breast cancer, compared to 91 percent in the control group.

Ninety-four percent of the women taking letrozole had survived at the five-year mark, compared to 93 percent of those taking a placebo.

In 2012, more than six million [women](#) around the world survived for at least five years after being diagnosed with breast cancer, according to statistics cited by ASCO.

© 2016 AFP

APA citation: Decade of hormone therapy reduces breast cancer recurrence: study (2016, June 5) retrieved 17 January 2021 from <https://medicalxpress.com/news/2016-06-decade-hormone-therapy-breast-cancer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.