


# Fighting penicilliosis—identification of fungal protein Mp1p as virulence factor

25 August 2016



Fungal cells (black arrows) in mouse liver infected with virulent *T. marneffeii* strain that produces high levels of Mp1p. Blue arrows indicate necrotic tissue. Credit: Woo et al. (2016)

Penicilliosis, caused by the fungus *Talaromyces marneffeii*, is a major opportunistic infection in Southeast Asia. A study published in *PLOS Neglected Tropical Diseases* reports that the fungal protein Mp1p is responsible for the pathogen's ability to cause disease in mice, and that it does so by allowing survival of the fungus in macrophages.

*T. marneffeii* infects bamboo rats, which are endemic in Southeast Asia. Humans who are immunocompromised because of HIV/AIDS or [immunosuppressive drugs](#) following [organ transplantation](#) are thought to be infected through exposure to bamboo rat excrements. Patients with penicilliosis have [poor prognosis](#) without treatment, and, even with treatment, mortality is approximately 20%.

Kwok-Yung Yuen, from the University of Hong

Kong, Hong Kong SAR, and colleagues have been working with *T. marneffeii*, a close relative of the *Penicillium* fungus, for decades. They had previously shown that a fungal protein called Mp1p, found on the pathogen's surface, can function like a vaccine—exposure to the protein induces protective immunity in mice.

To further study the role of Mp1p, the researchers started this study by searching the *T. marneffeii* genome for genes similar to the one encoding the Mp1p protein. The fungus, they found has 13 genes that are closely related to MP1. They named these relatives (or homologs) MPLP1-13. The researchers hypothesized that Mp1p and/or its homologs are so-called 'virulence factors' of *T. marneffeii*, which means they contribute to the pathogen's ability to cause disease.

To test this hypothesis, they systematically interfered with one of the 13 MPLP genes as well as MP1 itself. As the researchers were able to confirm, this generated *T. marneffeii* strains with dramatically reduced levels of individual Mp proteins. They then tested whether a fungal strain that produced no Mp1p or the 14 strains with low levels of individual Mp proteins could cause disease in mice. Sixty days after exposure to spores from the various strains, the researchers found that all mice infected with a strain that produced high levels of Mp1p died, even if they produced only low levels of one of the other Mp proteins. In contrast, mice infected with the strain that lacked Mp1p all survived. Infection with the strain that had low levels of Mp1p killed some but not all of the mice.

The survival data were consistent with different pathogen loads in mice infected with the different strains. Mice exposed to strains with high Mp1p had high level of fungus in their internal organs. Fungal levels in mice infected with the strain that lacked Mp1p were approximately 100-fold lower, and exposure to the strain with reduced Mp1p had

intermediate numbers of fungi in the examined tissues.

*T. marneffe*i Mp1p, the researchers found, was also able to turn a harmless fungus into a more virulent one. When the researchers introduced the *T. marneffe*i MP1 gene into the *Pichia* fungus (which doesn't have a related gene) and used the transgenic strain to infect mice, these mice had much higher levels of *Pichia* in their livers and spleens compared with mice infected with the parental *Pichia* strain.

Immune cells called macrophages (from Greek for 'big eaters') are the primary immune defense against fungal invaders such as *T. marneffe*i. To get at the mechanism by which Mp1p promotes virulence, the researchers tested the ability of [fungus](#) strains with different levels of Mp1p to survive after they are ingested by *T. marneffe*i macrophages. The more Mp1p, the researchers found, the higher the survival in macrophages.

Taken together, these results identify Mp1p as a bona fide virulence factor and potential drug target. As the researchers also discuss, *T. marneffe*i is closely related to the *Aspergillus* species—highly virulent molds that cause fatalities world-wide in patients with hematological malignancies, transplant recipients, HIV positive patients and patients on corticosteroid therapy. Several of the most dangerous *Aspergillus* fungi have proteins similar to Mp1p, and the researchers plan to investigate those in future work.

**More information:** Woo PCY, Lau SKP, Lau CCY, Tung ETK, Chong KTK, Yang F, et al. (2016) Mp1p Is a Virulence Factor in *Talaromyces* (*Penicillium*) *marneffe*i. *PLoS Negl Trop Dis* 10(8): e0004907. [DOI: 10.1371/journal.pntd.0004907](https://doi.org/10.1371/journal.pntd.0004907)

Provided by Public Library of Science

APA citation: Fighting penicilliosis—identification of fungal protein Mp1p as virulence factor (2016, August 25) retrieved 21 November 2019 from <https://medicalxpress.com/news/2016-08-penicilliosisidentification-fungal-protein-mp1p-virulence.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no*

*part may be reproduced without the written permission. The content is provided for information purposes only.*