


Study reveals protein to target in type 2 diabetes

1 September 2016, by Ziba Kashef



MARCH1 regulates insulin sensitivity by controlling cell surface insulin receptor levels, *Nature Communications* (2016). [DOI: 10.1038/ncomms12639](https://doi.org/10.1038/ncomms12639)

Provided by Yale University

Credit: stock.adobe.com

When the body's cells don't respond normally to insulin—a condition known as insulin resistance—blood glucose levels can increase, resulting in type 2 diabetes. Researchers have long known that insulin resistance is linked to defects in the insulin receptor (which controls glucose uptake) in multiple organs, including the liver.

To study the underlying mechanism, a team of researchers led by Narendra Wajapeyee, assistant professor of pathology, and Gerald Shulman, professor of cellular and molecular physiology and internal medicine, used a genomic technique to screen more than 600 proteins. They found that one of the proteins, MARCH1, impairs insulin by promoting the breakdown of the [insulin receptor](#) on the cell surface. MARCH1, which is increased in [obese individuals](#), could be a promising new target for drugs to treat type 2 diabetes, they said.

Lead authors on the study were Arvind Nagarajan and Max Petersen.

More information: Arvindhan Nagarajan et al.

APA citation: Study reveals protein to target in type 2 diabetes (2016, September 1) retrieved 23 June 2018 from <https://medicalxpress.com/news/2016-09-reveals-protein-diabetes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.