

# Marijuana use remains on the rise among US college students, but narcotic drug use declines

September 8 2016

---



A dried flower bud of the Cannabis plant. Credit: Public Domain

College student marijuana use continues its nearly decade-long increase,

according to the most recent national Monitoring the Future study.

In 2015, 38 percent of college students said they had used marijuana in the prior 12 months, up from 30 percent in 2006.

Daily or near-daily use of marijuana (having used 20 or more times in the prior 30 days) also has increased in recent years for college students, rising from 3.5 percent in 2007 to 5.9 percent in 2014—the highest level of daily use measured in the last 34 years.

However, in 2015 their daily use fell back some to 4.6 percent, or one in every 22 college students. A decline in the degree of risk of harm associated with using marijuana may account for much of the increase in use. Since 2003, proportions of 19-to-22-year-olds seeing regular use of marijuana as dangerous to the user has fallen sharply—from 58 percent in 2003 to 33 percent by 2015.

"This increase in use and decrease in perceived risk of harm regarding marijuana use should be taken seriously by college administrators, parents and students themselves. We know through other research that frequent marijuana use can adversely affect academic performance and college completion," said John Schulenberg, one of the study's lead researchers and research professor at the University of Michigan's Institute for Social Research.

In contrast to the story for marijuana use, other types of drug use are declining among college students. Nonmedical use of prescription narcotic drugs has been declining among college students since reaching a high in 2006 of 8.8 percent annual prevalence (that is, any use in the prior 12 months). By 2015, 3.3 percent of college students reported using any narcotic drug in the past 12 months without medical supervision—a drop of about six-tenths.

"It appears that college students, at least, are hearing and heeding the warnings about the very considerable dangers of using narcotic drugs," said Lloyd Johnston, the study's principal investigator and ISR distinguished senior research scientist and research professor.

Use of heroin, another narcotic drug, has been low among college students for many years. The highest annual prevalence recorded since 1980 was in 1998 at 0.6 percent, but the rate has been at or under 0.3 percent since 2005 and was down to 0.1 percent in 2015.

Use of amphetamines has started to decline among college students. From 2008 through 2012, the percentage of college students who reported using an amphetamine without medical supervision in the prior 12 months rose from 5.7 percent to 11.1 percent, likely due to more students using them to improve their [academic performance](#). But, by the time of the latest survey in 2015, that had fallen slightly to 9.7 percent.

"It appears that the increase in nonmedical use of prescription stimulant drugs may have passed its peak, though about one in 10 college students still report using them in the prior 12 months," Johnston said.

The use of MDMA (ecstasy and more recently "Molly") had made a bit of a comeback among college students between 2007 and 2012, but has been in decline since then. Annual prevalence in 2015 was 4.2 percent.

Certain drugs have declined in popularity quite rapidly among the nation's college students. For example, past-year use of synthetic marijuana, which is usually sold over the counter under such brand names as "K-2" and "Spice," dropped from 8.5 percent when first measured in 2011 to just 1.5 percent in 2015—a decline of about 80 percent. Salvia has fallen from 5.8 percent when its use was first measured in 2009 to just 0.4 percent in 2015—a decrease of more than 90 percent.

Some other drugs never gained much of a foothold on American college campuses. Past-year use of so-called "bath salts," a form of synthetic stimulants usually sold over the counter, has never exceeded 0.3 percent among college students since first being measured in 2012, and stands at 0.1 percent in 2015.

Past-year use of inhalants has been below 2 percent since 2005. The so-called "club drugs," Rohypnol and GHB, never really caught on, and have had negligible rates of use among college students.

Annual nonprescribed use of tranquilizers (4.3 percent) and sedatives (2.3 percent) has changed little in recent years, although these annual prevalence rates in 2015 are below those observed in the college population in the early 2000s.

In general, college males are more likely than college females to use nearly all of the illicit drugs. Sedatives are the primary exception; and there the genders are very close in their annual prevalence rates.

## **Cigarettes and Alcohol**

Cigarette smoking continues to decline gradually among college students, but the cumulative drop over the past 16 years has been dramatic. A peak rate of any smoking in the prior 30 days was reached in 1999 at 31 percent.

By 2015, the rate had fallen by nearly two-thirds to 11 percent, a record low. Daily smoking declined even more, from 19.3 percent in 1999 to 4.2 percent in 2015—a drop of nearly four-fifths and also a record low since 1980.

"The study shows that large declines in smoking rates have been occurring among secondary school students, as well," Johnston said. "So

much of the improvement among college students actually began when they were still in [high school](#)."

Their high school classmates not in college have dramatically higher rates of smoking. In 2015, 23 percent of them indicated past-month smoking, compared to 11 percent among the college students.

Heavy smoking is even more concentrated among those not in college, with their half-pack-or-more-per-day smoking rate at 9.1 percent versus 1.4 percent among college students.

"Cigarette smoking has become increasingly concentrated among the less educated," Johnston said.

Through 1993, college females had higher rates of smoking than college males; but since 1994, the opposite is true.

Electronic vaporizers, which include e-cigarettes, were used in the month prior to the survey by 14 percent of college males and 6 percent of college females.

Alcohol clearly remains the drug of choice among college students, with 79 percent indicating that they used in the past 12 months and 63 percent in the past 30 days.

Indeed, 62 percent said they were drunk at least once in the past 12 months and 38 percent in the past 30 days. So drinking and drunkenness remain quite commonplace on the nation's college campuses, even though there has been some modest falloff in these rates since the early 1980s.

Binge drinking—defined as having five or more drinks on at least one occasion in the past two weeks—was reported by 40 percent of all

college students in 2015 (close to the 38 percent who reported being drunk in the last month).

While the rate of binge drinking has gradually declined among college males over the past 30 years, there has been very little change in the rate among college females, resulting in some closing of the gap between genders (though males have consistently had a higher rate of binge drinking).

"Of even greater concern than binge drinking is what we have called 'extreme [binge drinking](#),' defined as having 10 or more drinks—or even 15 or more drinks—on at least one occasion in the prior two weeks," Johnston said.

Over the years 2011 to 2015 combined, about one in nine college students reported having 10 or more drinks in a row on at least one occasion in the prior two weeks, while one in 25 reported having 15 or more drinks in a row at least once in the same interval.

"Drinking at these levels can result in alcohol poisoning, serious accidents, and a host of unwise and dangerous behaviors," Johnston said. "So this remains a serious problem to be addressed."

The Monitoring the Future study is now in its 42nd year. Starting in 1980, the study has included nationally representative samples of full-time college students who are one to four years beyond high school. The annual samples of [college students](#) have ranged between 1,000 and 1,500 per year.

MTF is an investigator-initiated research undertaking, conceived and conducted by a team of research professors (listed as authors below) at the University of Michigan's Institute for Social Research. It is funded under a series of peer-reviewed, competitive research grants from the

National Institute on Drug Abuse, one of the National Institutes of Health.

MTF also conducts an annual national survey of high school seniors, from which a random, nationally representative subsample is drawn for follow-up by mail in future years. Of these follow-up respondents, those who are one to four years beyond high school and who report being in a two-year or four-year college full-time in March comprise the college student sample each year. They are not drawn from particular colleges or universities, which helps to make the sample more representative of the wide range of two- and four-year institutions of higher education in the country.

Provided by University of Michigan

Citation: Marijuana use remains on the rise among US college students, but narcotic drug use declines (2016, September 8) retrieved 26 April 2024 from <https://medicalxpress.com/news/2016-09-marijuana-college-students-narcotic-drug.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.