

# Kidney damage associated with imaging agent may be over-estimated

September 30 2016

---

A new analysis indicates that radiocontrast, which is commonly used during selected imaging tests may be less hazardous than previously thought. The findings, which appear in an upcoming issue of the *Journal of the American Society of Nephrology (JASN)*, suggest that imaging studies that might help save or improve lives are being unnecessarily withheld from patients owing to exaggerated fears.

While radiocontrast—sometimes referred to simply as "contrast" or "dye"—is frequently used to enhance imaging with computed tomography scans or [angiography](#), it can contribute to acute [kidney injury](#) (AKI). There is little agreement in the [medical literature](#) regarding the incidence of contrast-induced AKI, with published rates ranging from 30% of exposed patients.

To provide a more accurate estimate of the burden of AKI among patients receiving radiocontrast, Emilee Wilhelm-Leen, MD, Maria Montez-Rath, PhD, and Glenn Chertow, MD, MPH (Stanford University School of Medicine) analyzed 2009 information from the Nationwide Inpatient Sample, the largest publically available all-payer inpatient care database in the United States. After restricting the sample to hospitalizations for patients older than 18 years with lengths of stay  $\leq 10$  days, the investigators were left with 5,931,523 hospitalizations for analysis.

The researchers stratified patients according to the presence or absence of 12 relatively common diagnoses associated with AKI and evaluated

rates of AKI between strata. They also created a model that controlled for patients' health and comorbidities to estimate the risk of AKI associated with radiocontrast administration within each strata.

The team found that, in general, patients who received contrast did not develop AKI at a clinically significant higher rate than other patients. Patients who received radiocontrast developed AKI at a rate of 5.5%, compared with 5.6% of patients who did not receive radiocontrast. After controlling for patients' health and comorbidities, radiocontrast administration was actually linked to a 7% lower likelihood of developing AKI.

"We show data suggesting that the risk of acute kidney injury related to radiocontrast administration has been overestimated, and we would like for physicians, including cardiologists, radiologists, and surgeons who frequently are faced with decisions regarding the use or non-use of radiocontrast-enhanced imaging studies, to take this information into account in their clinical decision-making," said Dr. Chertow.

**More information:** E. Wilhelm-Leen et al. Estimating the Risk of Radiocontrast-Associated Nephropathy, *Journal of the American Society of Nephrology* (2016). [DOI: 10.1681/ASN.2016010021](https://doi.org/10.1681/ASN.2016010021)

Provided by American Society of Nephrology

Citation: Kidney damage associated with imaging agent may be over-estimated (2016, September 30) retrieved 26 April 2024 from <https://medicalxpress.com/news/2016-09-kidney-imaging-agent-over-estimated.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.