

# Adrenaline rush: Delaying epinephrine shots after cardiac arrest cuts survival rates

December 2 2016

---

Hospitals in which the administration of epinephrine to patients whose hearts have stopped is delayed beyond five minutes have significantly lower survival rates of those patients, a new study led by a cardiologist at UT Southwestern Medical Center finds.

Using data from a large registry, the national team of cardiologists found that nearly 13 percent of patients survived cardiac arrest when epinephrine shots were given within the first five minutes of the heart stopping, compared to about 11 percent when the epinephrine was given after five minutes, independent of all other aspects of care.

"That is a 20 percent better survival rate for patients at hospitals where epinephrine is given quickly, which is a big difference," said Dr. Rohan Khera, a Cardiology Division fellow at UT Southwestern and the first author on the study published online in the journal *Circulation*.

Delays in giving the epinephrine shots also had a negative effect on functional recovery, the researchers noted.

"These data are important for hospitals and patients. Improving epinephrine administration time, a likely correlate with overall CPR performance, may improve outcomes in cardiac arrest," said Dr. Mark Link, Professor of Internal Medicine at UT Southwestern and a specialist in heart-rhythm disorders, who was not involved in the study.

Researchers reviewed more than 100,000 records of patients whose heart

stopped while at the [hospital](#), using data from a large national registry run by the American Heart Association. The review involved nearly 550 hospitals across the country.

Researchers found considerable variability in how quickly epinephrine, commonly called adrenaline, was administered among the hospitals reviewed. Although individual hospitals were not identified in the study, the data showed that hospitals treating a high volume of cardiac arrest cases tended to administer the adrenaline shots more quickly than those with a lower volume of cases.

While TV shows commonly portray doctors dramatically shocking the heart with paddles, this is not the reality for most cardiac arrest [patients](#) in hospitals. About 80 percent of in-hospital cardiac arrests are due to causes that cannot be addressed using a defibrillator. These non-shockable heart stoppages are treated with CPR (cardio-pulmonary resuscitation) chest compressions and epinephrine, and these non-shockable cardiac arrests have much lower survival rates.

"Treatment options for non-shockable [cardiac arrest](#) are so limited that there has been an emphasis on improving current processes," said Dr. Khera. "Administering [epinephrine](#) promptly and improving the quality of CPR - these are the easily improved practices, which may be life-saving."

Researchers plan to examine processes at hospitals with few delays and compare processes at hospitals with more frequent delays to see if they can identify patterns that could help speed more timely administration of adrenaline shots and whether doing so would improve survival.

**More information:** Rohan Khera et al, Hospital Variation in Time to Epinephrine for Non-Shockable In-Hospital Cardiac Arrest, *Circulation* (2016). [DOI: 10.1161/CIRCULATIONAHA.116.025459](https://doi.org/10.1161/CIRCULATIONAHA.116.025459)

Provided by UT Southwestern Medical Center

Citation: Adrenaline rush: Delaying epinephrine shots after cardiac arrest cuts survival rates (2016, December 2) retrieved 26 April 2024 from <https://medicalxpress.com/news/2016-12-adrenaline-epinephrine-shots-cardiac-survival.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.