

Routine screening for genital herpes infection not recommended

December 20 2016

The U.S. Preventive Services Task Force (USPSTF) recommends against routine serologic screening (via a blood test) for genital herpes simplex virus infection in asymptomatic adolescents and adults, including those who are pregnant. The report appears in the December 20 issue of *JAMA*.

This is a D recommendation, indicating that there is moderate or high certainty that the screening has no net benefit or that the harms outweigh the benefits.

Genital herpes is a prevalent [sexually transmitted infection](#) in the United States; the Centers for Disease Control and Prevention estimates that almost 1 in 6 persons ages 14 to 49 years have genital herpes. Genital herpes [infection](#) is caused by 2 subtypes of [herpes simplex virus](#) (HSV), HSV-1 and HSV-2. Unlike other infections for which screening is recommended, HSV infection may not have a long asymptomatic period during which screening, early identification, and treatment may alter its course. Antiviral medications may provide symptomatic relief from outbreaks; however, these medications do not cure HSV infection. Although transmission of HSV can occur between an infected pregnant woman and her infant during vaginal delivery, interventions can help reduce transmission.

To update its 2005 recommendation on screening for genital herpes, the USPSTF reviewed the evidence on the accuracy, benefits, and harms of serologic screening for HSV-2 infection in asymptomatic persons,

including those who are pregnant, as well as the effectiveness and harms of preventive medications and behavioral counseling interventions to reduce future symptomatic episodes and transmission to others.

The USPSTF is an independent, volunteer panel of experts that makes recommendations about the effectiveness of specific preventive care services such as screenings, counseling services, and preventive medications.

Detection

In the past, most cases of genital herpes in the United States have been caused by infection with HSV-2. Adequate evidence suggests that the most widely used, currently available serologic screening test for HSV-2 approved by the U.S. Food and Drug Administration is not suitable for population-based screening, based on its low specificity, the lack of widely available confirmatory testing, and its high false-positive rate. Rates of genital herpes due to HSV-1 infection in the United States may be increasing. While HSV-1 infection can be identified by serologic tests, the tests cannot determine if the site of infection is oral or genital; thus, these serologic tests are not useful for screening for asymptomatic [genital herpes](#) resulting from HSV-1 infection.

Benefits of Early Detection and Intervention

Based on limited evidence from a small number of trials on the potential benefit of screening and interventions in asymptomatic populations and an understanding of the natural history and epidemiology of genital HSV infection, the USPSTF concluded that the evidence is adequate to bound the potential benefits of screening in asymptomatic adolescents and adults, including those who are pregnant, as no greater than small.

Harms of Early Detection and Intervention

Based on evidence on potential harms from a small number of trials, the high false-positive rate of the screening tests, and the potential anxiety and disruption of personal relationships related to diagnosis, the USPSTF found that the evidence is adequate to bound the potential harms of screening in asymptomatic adolescents and adults, including those who are pregnant, as at least moderate.

Summary

Based on the natural history of HSV infection, its epidemiology, and the available evidence on the accuracy of serologic screening tests, the USPSTF concluded that the harms outweigh the benefits of serologic [screening](#) for genital HSV infection in asymptomatic adolescents and adults, including those who are pregnant.

More information: *JAMA*, [DOI: 10.1001/jama.2016.16776](https://doi.org/10.1001/jama.2016.16776)

Provided by The JAMA Network Journals

Citation: Routine screening for genital herpes infection not recommended (2016, December 20) retrieved 19 September 2024 from <https://medicalxpress.com/news/2016-12-routine-screening-genital-herpes-infection.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.