

Head off 'holiday heart syndrome'

December 23 2016



(HealthDay)—Overindulging over the holidays can take a toll on the heart, according to experts at UT Southwestern Medical Center in Dallas.

Heavy drinking and eating too much, especially [salty foods](#), can trigger [atrial fibrillation](#). A-fib is an irregular heartbeat that causes the upper chambers of the heart to contract rapidly, increasing the risk for stroke.

People with the condition may feel short of breath, have [heart palpitations](#) and feel like their heart is beating much faster than usual.

During December when holiday revelry peaks, emergency rooms see a surge of patients with what's unofficially dubbed "holiday heart syndrome" from too much food and drink, according to UT Southwestern cardiologist Dr. Sharon Reimold.

"It's common for people to go to multiple parties during this time of year. You go to one party and have a drink or two, go to the next party and have a couple more. It's the cumulative effect of alcohol that can put you at risk, sending your heart into atrial fibrillation," Reimold said in a hospital news release.

Healthy adults should get no more than 1,500 milligrams of sodium a day, according to the American Heart Association. That's less than a teaspoonful. Reimold said it's important to watch what you're eating and drinking at holiday celebrations and avoid overdoing it.

More information: The American Heart Association provides more information on [holiday heart syndrome](#).

Copyright © 2016 [HealthDay](#). All rights reserved.

Citation: Head off 'holiday heart syndrome' (2016, December 23) retrieved 24 April 2024 from <https://medicalxpress.com/news/2016-12-holiday-heart-syndrome.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.