

Scientists identify two brain networks influencing how we make decisions

2 February 2017



Credit: Wikimedia Commons

Scientists at the Medical Research Council Brain Network Dynamics Unit at the University of Oxford have pinpointed two distinct mechanisms in the human brain that control the balance between speed and accuracy when making decisions.

Their discovery, published in *eLife*, sheds new light on the networks that determine how quickly we choose an option, and how much information we need to make that choice. A more detailed understanding of this intricate wiring in the [brain](#) holds the key to developing better treatments for neurological disorders such as Parkinson's disease.

The fundamental trade-off between speed and accuracy in [decision making](#) has been studied for more than a century, with a number of studies suggesting that the subthalamic nucleus region of the brain plays a key role.

"Previous behavioural studies of decision making do not tell us about the actual events or networks

that are responsible for making speed-accuracy adjustments," says senior author Peter Brown, Professor of Experimental Neurology at the University of Oxford. "We wanted to address this by measuring the exact location and timing of electrical activity in the subthalamic nucleus and comparing the results with behavioural data collected while a decision-making task is being performed."

Brown and his team first studied the reaction times of 11 patients with Parkinson's disease and 18 healthy participants, who were each asked to perform a moving-dots task. This required them to decide whether a cloud of moving dots appeared to be moving to the left or the right. The difficulty of the task was varied by changing the number of dots moving in one direction, and the participants were given randomly alternating instructions to perform the task with either speed or accuracy.

The researchers found that participants made much faster decisions when the task was easier - with the dots moving in a single direction - and when instructed to make a quick decision. They also found, in line with previous studies, that participants made significantly more errors during tests where they spent longer making a decision after being instructed to emphasise accuracy.

Using a computational model, they saw that it took longer in the more difficult tests for the brain to accumulate the necessary information to reach a critical threshold and make a decision. When the participants were asked to focus on speed, this threshold was significantly lower than when they focused on accuracy.

"The next step was to determine the activated networks in the brain that control these behavioural modifications and the trade-off between fast and accurate decisions," explains first author and postdoctoral fellow Damian Herz. "We measured the [electrical activity](#) of groups of nerve cells within

the subthalamic nucleus in patients with Parkinson's disease, who had recently been treated with [deep brain stimulation](#). We found two distinct neural networks that differ in the way they are ordered and the way they respond to tasks.

"One network increases the amount of information required before executing a decision and is therefore more likely to be activated when [accuracy](#) is important, while the second network tends to lower this threshold, especially when the choice needs to be made quickly."

The findings add to the increasing evidence that the pre-frontal cortex region of the brain contributes to decision making and opens up further interesting avenues to explore.

"We know that changes in activity of one of the sites we identified is also related to movement control," adds Brown. "Close relationships between these [neural networks](#) could mean that a common signal is responsible for adjustments in both the speed of decision and of the resulting movement. A better understanding of these mechanisms might make it possible to focus therapeutic interventions on specific neural circuits to improve treatment of neurological disorders in the future."

More information: Damian M Herz et al, Distinct mechanisms mediate speed-accuracy adjustments in cortico-subthalamic networks, *eLife* (2017). [DOI: 10.7554/eLife.21481](#)

Provided by eLife

APA citation: Scientists identify two brain networks influencing how we make decisions (2017, February 2) retrieved 25 September 2021 from <https://medicalxpress.com/news/2017-02-scientists-brain-networks-decisions.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.