


Clot removal therapy effective outside six-hour window for some stroke patients

May 16 2017



Micrograph showing cortical pseudolaminar necrosis, a finding seen in strokes on medical imaging and at autopsy. H&E-LFB stain. Credit: Nephron/Wikipedia

Results of a randomized controlled trial show that endovascular treatment (ET) to remove a stroke-causing blood clot in the brain is effective in some patients even when performed within 6 to 24 hours after a stroke. Current guidelines for acute stroke treatment endorse clot removal only when performed within six hours of a stroke. The findings

are presented today at the European Stroke Organisation Conference 2017 in Prague.

"This still means that you need to be rushed to the hospital as soon as possible after a [stroke](#) has occurred because the mantra "time is brain" still holds. However, our study shows that even if treated outside the time window, [patients](#) will have significantly reduced disability with clot removal," said Tudor Jovin, M.D., director, UPMC Stroke Institute, and professor of neurology and neurosurgery at the University of Pittsburgh, who co-led the trial.

The trial sets a new selection paradigm based on physiology, showing for the first time that looking at patients fulfilling certain imaging and clinical criteria, physiological state of the brain is a much better approach to determining whether patients will benefit from endovascular therapy as opposed to adhering to strict time windows, noted co-principal investigator Raul Nogueira, M.D., professor of neurology, neurosurgery and radiology at Emory University School of Medicine, and director of neuroendovascular service and neurocritical care service at Marcus Stroke & Neuroscience Center, Grady Memorial Hospital.

In the study, researchers randomly assigned stroke victims who arrived at the hospital outside the six-hour [time window](#) to either receive endovascular therapy or receive only standard medical therapy.

Brain imaging and clinical information (neurological deficit) was used to identify and enroll patients who had a small area of irreversibly damaged brain and a significantly larger [brain](#) tissue area that was imminently threatened by loss of blood supply, but still alive—a criterion known as clinical core mismatch.

The results showed that almost half of the patients (48.6 percent) receiving endovascular therapy had a good outcome at 90 days after

treatment—defined as the patients being independent in activities of daily living—showed clinical benefit while only 13.1 percent showed benefit in the group that received clot-busting drugs alone. There was no difference in mortality between the two groups.

The researchers planned to enroll 500 patients over the course of the study period. However, an intermediate review of the treatment effectiveness before enrollment was completed led the independent Data Safety Monitoring Board overseeing the study to recommend early termination of the trial after it was demonstrated that ET provided significant clinical benefit in the selected patients.

The multi-center international study known as the DAWN trial included trial locations in the United States, Spain, France, Australia and Canada. The trial was sponsored by Stryker Corporation, which manufactures the clot removal devices used in the study.

Provided by University of Pittsburgh Schools of the Health Sciences

Citation: Clot removal therapy effective outside six-hour window for some stroke patients (2017, May 16) retrieved 19 April 2024 from <https://medicalxpress.com/news/2017-05-clot-therapy-effective-six-hour-window.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--