

New consensus document for appropriate use of drug testing in clinical addiction medicine

June 1 2017

A new Consensus Document from the American Society of Addiction Medicine (ASAM) provides practical, evidence-based recommendations on the use of drug testing for identification, diagnosis, treatment, and monitoring of patients with or at risk for substance use disorders (SUDs). The document appears in the May/June issue of the *Journal of Addiction Medicine*, the official journal of ASAM.

"Drug testing is a valuable tool for supporting patients in [addiction](#) treatment, and this comprehensive set of recommendations should prove useful to providers in a variety of addiction treatment settings," comments Margaret Jarvis, MD, DFASAM, Chair of ASAM's Quality Improvement Council. The Consensus Document and supplemental digital content are available on [Journal of Addiction Medicine](#) and [ASAM](#) websites.

Evidence-Based Recommendations Aim at 'Smarter' Approach to Drug Testing

The Consensus Document provides practical guidance on the use of [drug testing](#) across the many clinical settings in which addiction treatment is provided. Based on evidence from more than 100 research studies, the document's recommendations were developed by a multidisciplinary panel, following an established "Appropriateness Method" process that combines scientific evidence with the collective judgment of experts.

The resulting recommendations focus on "when, where and how often it is appropriate to perform [drug](#) testing" in clinical [addiction medicine](#). Based on evidence that drug testing aids in monitoring adherence and abstinence in treatment and can improve patient outcomes, drug testing "should be widely used in addiction treatment settings." Drug testing should serve as a therapeutic tool, rather than a punitive one: "Providers should utilize drug testing to explore denial, motivation, and actual substance use behaviors with patients."

Drug testing plays important roles in initial assessment, treatment planning, and monitoring of patients with SUDs, across treatment settings. However, the document emphasizes, "Drug testing should be only one of several methods of detecting substance use or monitoring treatment," including patient self-reports and other indicators.

The Consensus Document addresses a wide range of daily issues encountered in the process of drug testing, including the choice of tests and appropriate response to test results. Urine tests remain by far the most common category of tests, but are also the most prone to tampering. There is growing interest in using other types of samples for drug testing, although more evidence on these tests is needed.

Drug testing should be done at least weekly at the beginning of treatment, and at least monthly in patients in stable recovery; however, individual consideration may be given for less frequent testing if a patient is in stable recovery. The document recommends that testing be performed on a random schedule, when possible.

The document addresses many other elements of testing programs, including documentation and patient confidentiality, practitioner education and expertise, and [test](#) facilities and devices. Issues in inpatient and outpatient treatment as well as opioid treatment services are discussed, along with issues related to special populations, including

adolescents, pregnant [patients](#), people in recovery, and health and other professionals. The expert panel also makes recommendations for further research to address gaps in each area.

A previous ASAM White Paper made broad suggestions to improve drug testing in clinical practice, emphasizing a "smarter" approach to drug testing—including avoiding some inappropriate and high-cost practices followed in the past. "The current crisis has many people in desperate need of high-quality addiction care, but many, professionals included, have no way of judging the quality of this kind of [treatment](#)," comments Dr. Jarvis. "This [document](#) helps to define that high-quality care."

More information: undefined undefined. Appropriate Use of Drug Testing in Clinical Addiction Medicine, *Journal of Addiction Medicine* (2017). [DOI: 10.1097/ADM.0000000000000322](https://doi.org/10.1097/ADM.0000000000000322)

Provided by Wolters Kluwer Health

Citation: New consensus document for appropriate use of drug testing in clinical addiction medicine (2017, June 1) retrieved 19 September 2024 from <https://medicalxpress.com/news/2017-06-consensus-document-drug-clinical-addiction.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--