

Artificial sweeteners linked to risk of weight gain, heart disease and other health issues

17 July 2017



Credit: CC0 Public Domain

Artificial sweeteners may be associated with long-term weight gain and increased risk of obesity, diabetes, high blood pressure and heart disease, according to a new study published in *CMAJ* (*Canadian Medical Association Journal*)

Consumption of artificial sweeteners, such as aspartame, sucralose and stevia, is widespread and increasing. Emerging data indicate that artificial, or nonnutritive, sweeteners may have negative effects on metabolism, [gut bacteria](#) and appetite, although the evidence is conflicting.

To better understand whether consuming artificial sweeteners is associated with negative long-term effects on weight and heart disease, researchers from the University of Manitoba's George & Fay Yee Centre for Healthcare Innovation conducted a systematic review of 37 studies that followed over 400 000 people for an average of 10 years. Only 7 of these studies were randomized controlled trials (the gold standard in clinical research), involving 1003 people followed for 6 months on average.

The trials did not show a consistent effect of artificial sweeteners on weight loss, and the longer observational studies showed a link between consumption of artificial sweeteners and relatively higher risks of weight gain and obesity, [high blood pressure](#), diabetes, [heart disease](#) and other health issues.

"Despite the fact that millions of individuals routinely consume artificial sweeteners, relatively few patients have been included in clinical trials of these products," said author Dr. Ryan Zarychanski, Assistant Professor, Rady Faculty of Health Sciences, University of Manitoba. "We found that data from clinical trials do not clearly support the intended benefits of artificial sweeteners for weight management."

"Caution is warranted until the long-term health effects of artificial sweeteners are fully characterized," said lead author Dr. Meghan Azad, Assistant Professor, Rady Faculty of Health Sciences, University of Manitoba. Her team at the Children's Hospital Research Institute of Manitoba is undertaking a [new study](#) to understand how artificial [sweetener](#) consumption by pregnant women may influence [weight gain](#), metabolism and gut bacteria in their infants.

"Given the widespread and increasing use of [artificial sweeteners](#), and the current epidemic of obesity and related diseases, more research is needed to determine the long-term risks and benefits of these products," said Azad.

The study was conducted by researchers from the University of Manitoba's George & Fay Yee Centre for Healthcare Innovation and the Children's Hospital Research Institute of Manitoba, Winnipeg, Manitoba.

More information: *Canadian Medical Association Journal* (2017).
www.cmaj.ca/lookup/doi/10.1503/cmaj.161390

Provided by Canadian Medical Association

Journal

APA citation: Artificial sweeteners linked to risk of weight gain, heart disease and other health issues (2017, July 17) retrieved 21 February 2018 from <https://medicalxpress.com/news/2017-07-artificial-sweeteners-linked-weight-gain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.