

Depression among young teens linked to cannabis use at 18

17 July 2017, by Bobbi Nodell



Cannabis indica. Credit: Wikipedia

A study looking at the cumulative effects of depression in youth, found that young people with chronic or severe forms of depression were at elevated risk for developing a problem with cannabis in later adolescence.

The study led by UW Medicine researchers interviewed 521 students recruited from four Seattle public middle schools. Researchers used data from annual assessments when students were ages 12-15 and then again when they were 18. The results were published in the journal *Addiction*.

"The findings suggest that if we can prevent or reduce chronic [depression](#) during early adolescence, we may reduce the prevalence of

cannabis use disorder," said lead author Isaac Rhew, research assistant professor of psychiatry and behavioral sciences at the University of Washington School of Medicine.

What researchers called "a 1 standard deviation increase" in cumulative depression during [early adolescence](#) was associated with a 50 percent higher likelihood of cannabis-use disorder.

According to researchers, during the past decade cannabis has surpassed tobacco with respect to prevalence of use among adolescents. Cannabis and alcohol are the two most commonly used substances among youth in the United States. They pointed to one national study showing increases in prevalence of cannabis use disorder and [alcohol use disorder](#) in the United States, especially among young adults.

Longitudinal studies looking at the link between depression and later use of alcohol and cannabis, however, have been mixed. Some show a link. Others don't. But most studies have assessed adolescent depression at a single point in time - not cumulatively, said the researchers. Further, there have been differences in how substance use has been measured ranging from the initiation of any use to heavier problematic forms of use.

The study oversampled for students with depressive and/or conduct problems. The researchers were surprised to see that the prevalence of cannabis and alcohol use disorder in this study was notably higher than national estimates with 21 percent meeting criteria for cannabis use disorder and 20 percent meeting criteria for alcohol use disorder at age 18.

What effect the easing of marijuana laws in Washington state had on the youth is unclear. Researchers said it would be informative to conduct a similar study in a state with more strict marijuana laws to understand whether the relationship

between depression and cannabis misuse would still hold in areas where marijuana may be less accessible.

The age 18 substance abuse assessments occurred between 2007-2010. Washington state legalized medical cannabis in 1998 and its medical cannabis market expanded greatly after 2009 when the federal justice department issued a ruling known as the Ogden Memo. And in 2003, the city of Seattle made [cannabis](#) offenses the lowest enforcement priority for police and the city attorney.

More information: Isaac C. Rhew et al. Examination of cumulative effects of early adolescent depression on cannabis and alcohol use disorder in late adolescence in a community-based cohort, *Addiction* (2017). DOI: [10.1111/add.13907](https://doi.org/10.1111/add.13907)

Provided by University of Washington

APA citation: Depression among young teens linked to cannabis use at 18 (2017, July 17) retrieved 26 September 2020 from <https://medicalxpress.com/news/2017-07-depression-young-teens-linked-cannabis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.