

# New PET-CT scan improves detection in rare cardiac condition

20 July 2017, by Jacqueline Carey

Using a new imaging technique that can diagnose cardiac sarcoidosis much more accurately than traditional tests, researchers from the University of Illinois at Chicago have found that the disease affects other organs in 40 percent of patients with cardiac sarcoidosis. The findings are published in the *Journal of Nuclear Cardiology*.

Sarcoidosis is an inflammatory disease that often attacks multiple organs, particularly the lungs, heart and lymph nodes, and disproportionately affects African-Americans. Sarcoidosis can be very difficult to diagnose using available tests and available imaging techniques.

Positron emission tomography (PET) scans, and computed tomography (CT) scans, as well as scans that combine information from each, have been used in the diagnosis of cardiac sarcoidosis, but the results are often unclear and hard to interpret, and lead to many false negatives.

A new combined PET-CT scan [technique](#) developed at UIC incorporates a 72-hour high-fat, low-sugar diet before images are taken. The new scan provides much clearer images of sarcoidosis and lets physicians better-detect cardiac sarcoidosis. In a previous UIC study, researchers were able to diagnose cardiac sarcoidosis much more accurately and could diagnose cases of cardiac sarcoidosis that would not have been picked up using the usual 24-hour high-fat, low-sugar protocol normally prescribed before the combined PET-CT scans.

In the current study, the researchers used the new protocol to investigate the relationship between the presence of cardiac sarcoidosis and sarcoidosis in other parts of the body.

"We wanted to know if patients being evaluated for cardiac sarcoidosis would benefit from more thorough imaging of the body beyond the usual torso scan, using this new technique," said Dr.

Nadera Sweiss, professor of rheumatology in the UIC College of Medicine and an author on the paper.

The researchers performed full body (nose to knees) PET-CT scans of 188 patients who followed the 72-hour pre-test diet between Dec. 2014 and Dec. 2015 at the University of Illinois hospital, UI Health. Of the 20 scans that were positive for cardiac sarcoidosis, eight, or 40 percent, revealed sarcoidosis in other parts of the body.

"This study indicates that more extensive, full-body scans of patients with known cardiac sarcoidosis can uncover secondary sites of the disease in a significant number of patients," said Sweiss, who is also director of UI Health's Bernie Mac Sarcoidosis Translational Advanced Research (STAR) Center. "Knowing there is disease in organs other than the heart changes the way we approach treatment—it allows us to more accurately stage the [disease](#) and treat it accordingly."

Sweiss cautions that the technique for detecting sarcoidosis outside and within the heart needs to be further tested, and its ultimate contribution to outcomes in [patients](#) with cardiac sarcoidosis will need to be evaluated.

**More information:** Darshan C. Patel et al, FDG PET-CT findings of extra-thoracic sarcoid are associated with cardiac sarcoid: A rationale for using FGD PET-CT for cardiac sarcoid evaluation, *Journal of Nuclear Cardiology* (2017). [DOI: 10.1007/s12350-017-0962-4](#)

Provided by University of Illinois at Chicago

APA citation: New PET-CT scan improves detection in rare cardiac condition (2017, July 20) retrieved 3 March 2021 from <https://medicalxpress.com/news/2017-07-pet-ct-scan-rare-cardiac-condition.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*