


Researchers uncover how to boost learning efficiency in neurofeedback

July 31 2017



Credit: National Research University Higher School of Economics

Researchers from the HSE Centre for Cognition & Decision Making and the (Institute of Problems of Mechanical Engineering, Russian Academy of Sciences, have conducted a series of experiments to determine what a

person actually controls when tasked with independently affecting the activity of their own brain. This discovery may contribute to non-pharmacological methods for treating epilepsy, attention-deficit/hyperactivity disorder and depression. The research results were published in the journal *Scientific Reports*.

The human [brain](#) is always electrically active. Neurons exchange information using very short electrical impulses. The aggregate of a large number of these impulses, which are distributed in time and appear in various parts of the brain, cause fluctuations in the electrical potential of the head's surface. Similar to a cardiogram, we can use methods such as electroencephalograms to register these fluctuations and judge brain activity. Further, several types of activity can be determined by the speed of the fluctuations: alpha, beta, gamma, theta, and delta waves. With electroencephalography, these waves can be registered, recoded into a perceivable form such as the brightness of a computer screen, and presented in real time. This serves as the basis for neurofeedback technology, which is one type of biofeedback. With the help of biofeedback, a person can learn to regulate the activity of his or her own brain.

The first time neurofeedback was discussed as a method for teaching a person to regulate the electromagnetic activity of their brain was in research conducted by Joseph Kamiya of the University of Chicago in the 1960s. Kamiya showed that in receiving feedback on brain activity, a person learns to affect this activity while they learn to determine the current state of their [brain activity](#) and, for example, the prevalence of a particular kind of rhythm.

In the experiment, the researchers used electroencephalography to assess the intensity of the brain's alpha wave activity. Alpha waves relate to a person's ability to relax, manage stress, and internalise new information. The research subjects saw information about their alpha wave activity in

the form of red colour saturation on a monitor and tried to mentally make this colour more saturated.

The study involved 18 people, nine of whom (the experimental group) received real feedback on their efforts, i.e., they saw the colour's saturation reflecting the current level of alpha wave activity. Conversely, the control group received false feedback, meaning changes in the intensity of the colour displayed on the screen were not connected with the research subjects' alpha wave activity.

Over the course of two days, both groups carried out five two-minute sessions per day. The research subjects were not provided with a strategy that would produce the best results.

At the end of the first day, the entire experimental group showed an increase in alpha wave activity, while alpha wave activity was slightly lower at the beginning of the second day than at the end of the first. But throughout the second day, activity continued to grow and exceeded the levels seen at the end of the first day. Two days of training in the neurofeedback paradigm were enough to significantly increase alpha wave capacity. This sort of increase was not seen in the control group.

Alpha wave activity is non-stationary and occurs in the form of special bursts, each of which can be characterised by its duration and amplitude. Another important parameter is the number of such bursts per unit of time.

"We became interested in what exactly changes during feedback—the amplitude of each burst, its length, or how often these bursts occur," explains Professor Alexey Ossadtchi, who is a senior research fellow in the Centre for Cognition & Decision Making and one of the study's authors.

"These are three completely different metrics as far as neurophysiology is concerned. When amplitude increases, the size of the corresponding neuron population increases, and when the duration grows, short-term regulatory mechanisms with feedback are directly involved and allow the brain to maintain high alpha wave activity for a longer period of time. An increase in the frequency of bursts shows that under the influence of a person's intentional effort, the brain more easily enters a state in which the alpha rhythm dominates."

After the experimenters compared how much the amplitude, duration, and frequency of bursts changed during the training process of all participants on different days, it was discovered that alpha wave bursts in particular undergo the most significant changes per unit of time. In other words, by training in the neurofeedback paradigm, a person learns to control entry into a completely alpha state. This is new information in the study of neurofeedback mechanisms.

"It turned out that the frequency of alpha wave bursts in particular is a coached metric that we can use to influence alpha wave activity as a whole, unlike amplitude and duration, which are likely coded at a lower level," Professor Ossadtchi adds. "This means that in therapy and in training, we have to give people feedback specifically on the parameter that they are truly able to influence—in our case, for every entry into the [alpha] state and for every new burst. It's preferred that a person receive such reinforcement with as little delay as possible. That is what we are working on now."

The researchers assume that the data they collected are valid not only for [alpha waves](#), but for other types of electromagnetic frequencies, as well. A similar experiment will be conducted soon on different types of waves.

The neurofeedback training paradigm lowers the likelihood of epileptic

seizures, eliminates some manifestations of attention deficit/hyperactivity disorder, and provides relief to a person with depression. In addition, neurofeedback as a technology helps athletes control their psycho-emotional state, and it can also be used to master the art of meditation, improve memory, and increase a person's ability to concentrate. According to the researchers, knowing specifically which regulatory mechanisms are activated during a certain form of neurofeedback allows one to significantly increase the efficiency of this technology, while it also provides access to new resources in the [human brain](#) that have not yet been studied.

More information: Alexei Ossadtchi et al, Neurofeedback learning modifies the incidence rate of alpha spindles, but not their duration and amplitude, *Scientific Reports* (2017). [DOI: 10.1038/s41598-017-04012-0](https://doi.org/10.1038/s41598-017-04012-0)

Provided by National Research University Higher School of Economics

Citation: Researchers uncover how to boost learning efficiency in neurofeedback (2017, July 31) retrieved 26 April 2024 from <https://medicalxpress.com/news/2017-07-uncover-boost-efficiency-neurofeedback.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--