

Patient profile impacts QOL with radiation Tx in head, neck cancer

11 August 2017



"Clinicians can use knowledge on time course and associated characteristics to identify and inform patients at higher risk for diminished QOL," the authors write.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2017 [HealthDay](#). All rights reserved.

(HealthDay)—There are patient characteristics and clinical factors independently associated with physical and mental quality of life (QOL) in patients with head and neck cancer undergoing radiotherapy, according to a study published online Aug. 2 in *Head & Neck*.

Guro Lindviksmoen Astrup, from Oslo University Hospital in Norway, and colleagues evaluated 133 patients with head and [neck cancer](#). From the initiation of radiotherapy through six months of follow-up, patients rated physical and mental QOL using the Short Form-12 five times.

The researchers found that QOL deteriorated during radiotherapy and gradually improved after completion. There was a negative association between less [social support](#) and both physical and mental QOL. There were also negative associations between older age, more comorbidities, more psychological symptoms, and concomitant chemotherapy with physical QOL. There were positive associations for male sex, fewer physical symptoms, surgery before [radiotherapy](#), and concomitant chemotherapy with mental QOL.

APA citation: Patient profile impacts QOL with radiation Tx in head, neck cancer (2017, August 11) retrieved 18 October 2017 from <https://medicalxpress.com/news/2017-08-patient-profile-impacts-qol-tx.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.