


# Memory decline after head injury may be prevented by slowing brain cell growth

15 September 2017



Credit: copyright American Heart Association

The excessive burst of new brain cells after a traumatic head injury that scientists have traditionally believed helped in recovery could instead lead to epileptic seizures and long-term cognitive decline, according to a new Rutgers New Jersey Medical School study.

In the September issue of *Stem Cell Reports*, Viji Santhakumar, associate professor in the department of Pharmacology, Physiology and Neuroscience, and her colleagues, challenge the prevailing assumption by scientists in the field that excessive neurogenesis (the birth of new brain cells) after [injury](#) is advantageous.

"There is an initial increase in birth of new neurons after a brain injury but within weeks, there is a dramatic decrease in the normal rate at which neurons are born, depleting brain cells that under

normal circumstances should be there to replace damaged cells and repair the brain's network," said Santhakumar. "The excess new neurons lead to [epileptic seizures](#) and could contribute to [cognitive decline](#)"

In the United States an estimated 1.7million people sustain a TBI each year, making the condition a major cause of death and disability. Symptoms can include impaired thinking or memory, personality changes and depression and vision and hearing problems as well as epilepsy. About 80 percent of those who develop epilepsy after a brain injury have seizures within the first two years after the damage occurs.

Santhakumar said while researchers who study epilepsy have started to look more closely at how preventing excessive neurogenesis after brain injury could prevent seizures, neuroscientists have traditionally viewed the process as helpful to overall brain recovery.

Studying laboratory rats, Rutgers scientists found, however, that within a month after experimental brain injury, the number of new brain cells declined dramatically, below the numbers of new neurons that would have been detected if an injury had not occurred.

When scientists were able to prevent the excessive neurogenesis which occurs within days of the injury with a drug similar to one under trial for chemotherapy treatments, the rate of birth of new brain cells went back to normal levels and risk for seizures was reduced.

"That's why we believe that limiting this process might be beneficial to stopping seizures after [brain injury](#)," she said.

While the regenerative capability of brain [cells](#), in the hippocampus - the part of the [brain](#) responsible for learning and memory - slows down as part of

the aging process, the Rutgers scientists determined that the process that occurred after a head injury was related to injury and not age.

"It is normal for the birth of new neurons to decline as we age," said Santhakumar. "But what we found in our study was that after a head injury the decline seems to be more rapid."

Provided by Rutgers University

APA citation: Memory decline after head injury may be prevented by slowing brain cell growth (2017, September 15) retrieved 27 January 2022 from <https://medicalxpress.com/news/2017-09-memory-decline-injury-brain-cell.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*