

One weight-loss surgery shows lasting results

September 21 2017, by Amy Norton, Healthday Reporter



(HealthDay)—Obesity surgery can have long-lasting effects on weight and the risk of developing type 2 diabetes and high blood pressure, a new study finds.

Doctors have known that a type of obesity [surgery](#), called gastric bypass, works in the short-term. Patients typically lose a lot of weight, and obesity-related health problems can be prevented or even cured.

But the new findings show that the benefits are still apparent 12 years later.

The study, of more than 1,100 severely obese adults, found that those who underwent gastric bypass lost an average of 100 pounds over two years. By year 12, they'd managed to keep 77 of those pounds off.

On top of the weight loss, [surgery patients](#) had a much lower risk of developing type 2 diabetes—92 percent lower, versus obese patients who did not have surgery.

"This is very effective at diabetes prevention," said lead researcher Ted Adams, a professor at the University of Utah School of Medicine.

The surgery can also reverse existing diabetes. At year 12, half of patients who'd had type 2 diabetes before surgery were in remission, according to the study.

Still, Adams said, gastric bypass has risks, and it's considered a last resort after other weight-loss attempts have failed.

So candidates for the surgery should go in fully informed of the pros and cons, Adams said.

An obesity specialist who was not involved in the study agreed.

"It does take a lot of [hard work](#) before and after surgery. And this is not a cure-all for diabetes," said Dr. Scott Isaacs, medical director of Atlanta Endocrine Associates.

The hard work includes a lifelong commitment to diet and exercise changes. So the surgery itself is no "magic bullet," said Isaacs, who is a spokesperson for the Obesity Society.

That said, he called the news findings "really clear-cut."

"It's one of the best studies we've had done to date," Isaacs said. "And it shows that this is a safe, effective surgery with durable results."

The study, funded by the U.S. National Institutes of Health (NIH), involved 1,156 severely obese adults.

In all, 418 underwent gastric bypass, while the rest either did not seek surgery or considered it but did not go through with it—mostly for insurance reasons.

During [gastric bypass](#), a surgeon staples the stomach to create a "pouch" that can hold only a small amount of food at a time. Then a passage is created from the pouch to the middle of the small intestine—limiting the body's absorption of nutrients.

In general, Isaacs said, the surgery is reserved for people with a [body mass index](#) (BMI) of at least 40 —meaning they're around 100 pounds or more overweight. It may also be recommended for people who have a BMI of 35-plus and health conditions such as diabetes or heart disease.

There are risks from the surgery itself, including blood clots, infections and bleeding, according to the NIH.

"It's a major surgery," Adams said, "and you can't reverse it."

The risk of death, he added, is low—on par with hip replacement surgery—but it's a possibility.

After surgery, Isaacs said, there's a risk of nutrient deficiencies, so patients need to take prescribed supplements.

All of that has to be weighed against the potential benefits, Adams said.

Of surgery patients in his study, 3 percent developed diabetes over the next dozen years. That compared with 26 percent of nonsurgery patients.

Similarly, 16 percent developed [high blood pressure](#), versus more than 40 percent of other patients.

Of patients who'd had [diabetes](#) before surgery, 75 percent went into remission by year two. That waned to 51 percent by year 12.

Still, Adams said, the long-term remission rate is "pretty remarkable."

There was one concerning finding—one that past studies have uncovered, too. Seven study patients died by suicide after having surgery.

It's not clear why, Adams said. But [patients](#) in the surgery group did tend to report a poorer quality of life before having the procedure, he said, versus people who did not opt for surgery.

It's possible, Adams speculated, that for certain people, the surgery did not improve their life to the degree they'd hoped.

Also, Isaacs noted that research suggests the surgery may change the way alcohol is metabolized—and potentially contribute to drinking problems in certain people.

The study findings were published in the Sept. 21 issue of the *New England Journal of Medicine*.

More information: Ted Adams, Ph.D., M.P.H., Intermountain Healthcare Live Well Center, and adjunct professor, internal medicine, University of Utah School of Medicine, Salt Lake City; Scott Isaacs, M.D., endocrinologist and medical director, Atlanta Endocrine Associates, Atlanta, and spokesperson, The Obesity Society, Silver Spring, Md.; Sept. 21, 2017, *New England Journal of Medicine*

The U.S. National Institute of Diabetes and Digestive and Kidney Diseases has more on [obesity surgery](#).

Copyright © 2017 [HealthDay](#). All rights reserved.

Citation: One weight-loss surgery shows lasting results (2017, September 21) retrieved 20 September 2024 from <https://medicalxpress.com/news/2017-09-weight-loss-surgery-results.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--