

Arterial stiffness linked to incidence of diabetes

29 October 2017



"Increased c-f PWV is associated with increased incidence of diabetes, independent of other risk factors. These results suggest that increased arterial stiffness is an early risk marker for developing [diabetes](#)," the authors write.

More information: [Abstract/Full Text](#) (subscription or payment may be required)

Copyright © 2017 [HealthDay](#). All rights reserved.

(HealthDay)—Increased arterial stiffness, as measured by carotid-femoral pulse wave velocity (c-f PWV), is associated with increased incidence of diabetes, according to a study published online Sept. 29 in *Diabetes Care*.

Iram Faqir Muhammad, from Lund University in Sweden, and colleagues examined the correlation between [arterial stiffness](#) as measured by c-f PWV and incidence of diabetes in a population of participants from the Malmö Diet and Cancer cardiovascular cohort. Data were included for 2,450 individuals (mean age, 71.9±5.6 years).

The researchers found that 68 participants (2.8 percent) developed diabetes during a mean follow-up of 4.43±1.4 years. Subjects in the first, second, and third tertiles of c-f PWV had crude incidence of diabetes of 3.5, 5.7, and 9.5, respectively, per 1,000 person-years. The hazard ratio for diabetes was 1 (reference), 1.83, and 3.24, respectively, for the tertiles of c-f PWV after adjustment for potential confounders.

APA citation: Arterial stiffness linked to incidence of diabetes (2017, October 29) retrieved 27 September 2020 from <https://medicalxpress.com/news/2017-10-arterial-stiffness-linked-incidence-diabetes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.