

Parents help shape how much pain preschoolers feel after vaccination

14 November 2017



York University Professor Rebecca Pillai Riddell in the Faculty of Health, York Research Chair in Pain and Mental Health and senior author of the paper and her team found that not only is a parent's behaviour during vaccinations critical to a child's pain coping responses, but that the behaviour may also impact their reactions in the future. Moreover, the research may better inform medical care and may predict suffering by children during vaccinations into adulthood. Credit: York University

While vaccinations protect children against various illnesses, the pain can sometimes be too much to bear. It's no wonder most children and parents dread their vaccination appointments. Now new research from [York University's OUCH Cohort](#) at the Faculty of Health found that the amount of distress and pain felt by a preschooler during a vaccination is strongly related to how their parents help them cope before and during an appointment.

Professor Rebecca Pillai Riddell in the Faculty of Health, York Research Chair in Pain and Mental Health and senior author of the paper, has been following the OUCH Cohort [children](#) for over a decade. In the study, researchers used the data from 548 children who had been followed during infant and/or preschool vaccinations. Infants were included in the study if the infant had no suspected

developmental delays or impairments, had no chronic illnesses, had never been admitted to a [neonatal intensive care](#) unit, and was born no more than three weeks preterm.

The research, led by graduate student Lauren Campbell, examined children who were expressing the most pain during preschool vaccinations. The goal of the study was to find out what would best predict the children who had the highest pain and did the poorest coping during the preschool vaccination by watching both the child and the parent over repeated vaccinations over childhood. Researchers evaluated various pain behaviours such as facial activity (grimacing), leg activity (crunching of legs), crying and consolability to measure the level of pain in children. They also looked at what the child and parent said that related to coping with the pain.

The results suggested that a preschooler's ability to cope is a powerful tool to reduce pain-related distress but they need [parents](#) to support their coping throughout a vaccination appointment to have an impact in reducing pain-related distress.

"When children were distressed prior to the needle, that made them feel more pain after the needle," says Pillai Riddell.

The data confirmed that engaging in coping-promoting behaviours like encouraging a child to take deep breaths was important. Using distractions such as pulling out an iPhone or distracting children with plans about what they will do after the appointment also improved children's coping.

However, Pillai Riddell says it may be even more important to avoid negative or distress-promoting behaviours.

"Telling kids that 'it's ok, it's going to be fine' over and over again actually makes children feel

anxious. Parents only say things are 'okay' when things are not ok. Ensuring you don't criticize a child, such as saying: 'strong girls don't cry', 'big boys don't do that' is important. Also, don't apologize to a child by saying things like: 'I'm sorry this is happening to you,' is also key, says Pillai Riddell. "These are all distress-promoting behaviours and increase pain and distress."

The study, published in *Pain*, found that not only is a parent's [behaviour](#) during vaccinations critical to a child's [pain](#) coping responses, but that the behaviour may also impact their reactions in the future. Moreover, the research may better inform medical care and may predict suffering by children during vaccinations into adulthood.

"People who have negative reactions with doctors when they are young, may avoid preventative care in the future. If you didn't like a needle when you were five, that can stick with you."

More information: Lauren Campbell et al, Preschool children's coping responses and outcomes in the vaccination context, *PAIN* (2017).
[DOI: 10.1097/j.pain.0000000000001092](https://doi.org/10.1097/j.pain.0000000000001092)

Provided by York University

APA citation: Parents help shape how much pain preschoolers feel after vaccination (2017, November 14) retrieved 20 June 2018 from <https://medicalxpress.com/news/2017-11-parents-pain-preschoolers-vaccination.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.