

Aspirin can prevent heart attacks after noncardiac surgery in patients with prior PCI

14 November 2017



Graham found that aspirin can help prevent heart attacks in non-cardiac surgery patients who previously had a coronary procedure such as an angioplasty or stent. Credit: Melissa Fabrizio

A Canadian-led study has found that perioperative aspirin can prevent heart-related complications after major noncardiac surgery in patients with previous percutaneous coronary intervention (PCI) such as an angioplasty or stent.

The study found that for every 1,000 [patients](#) with PCI, giving them [aspirin](#) at the time of [noncardiac surgery](#) would prevent 59 heart attacks and cause eight major bleeding events. The results of this study are significant, given that 200 million adults undergo major noncardiac surgery annually.

These results were a substudy of POISE-2, a large international study with sites in 135 centres in 23 countries, including Edmonton. Patients with previous PCI were enrolled in 82 centres in 21 countries.

Noncardiac surgeries occur daily at hospitals

around the world, so the study results will have a big impact on this patient group.

"This is your next-door neighbour who had angioplasty five years ago, feels fine and needs to go in for hip surgery. It affects quite a large number of people," said Michelle Graham, an interventional cardiologist, professor in the University of Alberta's Department of Medicine and lead of the substudy. "We believe that perioperative aspirin in this group of patients will most likely benefit them and reduce their risk of serious vascular complications."

In patients without a PCI, POISE-2 found that aspirin did not reduce the risk of heart attack and led to an increased risk of major bleeding.

Of the 10,010 participants enrolled in the POISE-2, 470 had a previous PCI. Since patients with a prior PCI have an increased risk of cardiovascular complications after noncardiac surgery, the group wanted to see if the findings were the same in the subgroup.

This was the largest randomized trial of patients with PCI undergoing major noncardiac surgery. Although the study found that aspirin reduced the risk of [heart attack](#), it did slightly increase the risk of bleeding, though it did not appear worse than in the overall POISE-2 trial.

"There will be a big knowledge translation push with our colleagues in anesthesia and [surgery](#) to remind them we want them to continue aspirin in this group of patients, when for most other groups we're recommending they stop," says Graham.

The Canadian Cardiovascular Society guidelines published last year indicated that aspirin should not be given in the perioperative period.

"This will potentially change the practice of anyone who does perioperative medicine," said Graham.

Graham recognizes that studies like this are not possible without the participation of patients.

"We could not advance the field of medicine without the patients that participate in trials like POISE-2," said Graham. "This really is a team sport, we need the surgeons, anesthesiologist and patients on board to make it successful."

The study 'Aspirin in Patients with Previous Percutaneous Coronary Intervention Undergoing Noncardiac Surgery' was published in *Annals of Internal Medicine* on November 14, 2017.

More information: *Annals of Internal Medicine* (2017). DOI: [10.7326/M17-2341](https://doi.org/10.7326/M17-2341)

Provided by University of Alberta Faculty of
Medicine & Dentistry

APA citation: Aspirin can prevent heart attacks after noncardiac surgery in patients with prior PCI (2017, November 14) retrieved 18 February 2018 from <https://medicalxpress.com/news/2017-11-aspirin-heart-noncardiac-surgery-patients.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.