

Popular blood pressure medicine linked with increased risk of skin cancer

5 December 2017

Recently published research from The University of Southern Denmark and the Danish Cancer Society shows a connection between one of the most common medications for hypertension and skin cancer. Danish researchers set their sights on anti-hypertensive medicine containing hydrochlorothiazide in relation to an increased risk for skin cancer. The researchers have previously demonstrated that the medicine, which is one of the most commonly used drugs worldwide, can increase the risk of lip cancer. medicine is now associated, can be easily treated and has a very low mortality rate.

In a new study, the researchers have identified another clear connection between the use of hypertension [medicine](#) and the chance of developing [skin cancer](#). More specifically, this refers to drugs containing hydrochlorothiazide and [squamous cell carcinoma](#).

Surprisingly high risk

The researchers have also looked at other commonly used hypertension medicines, but none of them increased the risk of [skin](#) cancer.

"We knew that hydrochlorothiazide made the skin more vulnerable to damage from the sun's UV rays, but what is new and also surprising is that long-term use of this [blood pressure](#) medicine leads to such a significant increase in the risk of skin cancer," says Anton Pottegård, Associate Professor, Ph.D., from the University of Southern Denmark.

The study, which is based on about 80,000 Danish cases of skin cancer, shows that the risk of developing skin cancer is up to seven times greater for users of medicine containing hydrochlorothiazide.

Risk of impairment

There are several types of skin cancer. Squamous cell carcinoma, with which the blood pressure

"However, both lip and skin cancer are typically treated with an operation that is associated with a certain risk of impairment as well as a small but real risk that skin cancer of the squamous cell type spreads," says Anton Pottegård.

Hydrochlorothiazide is one of the most commonly used medicines to reduce blood pressure both in the U.S., where over 10 million people use the drug annually, and in Western Europe. The side effects therefore affect many people, and the researchers have calculated that about 10 percent of all Danish cases of squamous cell carcinoma may be caused by hydrochlorothiazide.

"You should not interrupt your treatment without first consulting your doctor. However, if you use hydrochlorothiazide at present, it may be a good idea to talk to your doctor to see if it is possible to choose a different medicine," says Anton Pottegård.

As part of the project, the Danish researchers partnered with Dr. Armand B. Coggnetta Jr., Chief Division of Dermatology, Florida State University, who is the head of one of the largest treatment centres for skin cancer in the US. Dr. Coggnetta has extensive experience in diagnosing and treating skin cancer patients in Florida, and he has noted that hydrochlorothiazide is suspiciously prevalent among his patients, particularly among so-called "catastrophic patients," who can have several hundred skin cancers each. He welcomes the new results.

"We have seen and followed many patients with different skin cancers where the only risk factor apart from exposure to sunlight seems to be hydrochlorothiazide. The combination of living and residing in sunny Florida while taking hydrochlorothiazide seems to be very serious and

even life-threatening for some patients. Even though we knew that hydrochlorothiazide made you sensitive to the sun, the connection between this medicine and skin cancer has remained elusive. The study carried out by Pottegård and his colleagues will have great impact on [skin cancer prevention](#) and public health worldwide," he explains.

What now?

The researchers continue to work on studies that can shed additional light on the connection between hydrochlorothiazide and skin cancer. Furthermore, they have entered into a dialogue with relevant medical companies as well as the Danish Medicines Agency regarding their findings.

"The [risk](#) of skin [cancer](#) must, of course, be weighed against the fact that hydrochlorothiazide is an effective and otherwise safe treatment for most patients. Nevertheless, our results should lead to a reconsideration of the use of hydrochlorothiazide. Hopefully, with this study, we can contribute towards ensuring safer treatment of high blood pressure in the future," says Anton Pottegård.

More information: [DOI: 10.1016/j.jaad.2017.11.042](#)

Provided by University of Southern Denmark

APA citation: Popular blood pressure medicine linked with increased risk of skin cancer (2017, December 5) retrieved 25 June 2019 from <https://medicalxpress.com/news/2017-12-popular-blood-pressure-medicine-linked.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.