

# Running away from addiction: How exercise aids smoking cessation

20 December 2017

(2017). [DOI: 10.1111/bph.14068](https://doi.org/10.1111/bph.14068)



Provided by Wiley

Credit: CC0 Public Domain

New research in mice sheds light on the mechanism underlining exercise's protective effect against nicotine dependence and withdrawal.

The *British Journal of Pharmacology* study reveals that exercise during [nicotine exposure](#) markedly reduces the severity of nicotine withdrawal symptoms, an effect that is accompanied by increased activation of  $\alpha 7$  nicotinic acetylcholine receptors (which are targets of nicotine) in the [hippocampal region](#) of the brain.

"The findings support the protective effect of exercise preceding smoking cessation against the development of physical dependence, which may aid smoking cessation by reducing the severity of withdrawal symptoms," said senior author Dr. Alexis Bailey, of St. George's University of London.

**More information:** Helen Keyworth et al, Wheel running during chronic nicotine exposure is protective against mecamylamine-precipitated withdrawal and upregulates hippocampal  $\alpha 7$  nACh receptors in mice, *British Journal of Pharmacology*

APA citation: Running away from addiction: How exercise aids smoking cessation (2017, December 20) retrieved 16 September 2021 from <https://medicalxpress.com/news/2017-12-addiction-aids-cessation.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*